



Investigating the Eastern Alpine–Dinaric transition with teleseismic receiver functions: Evidence for subducted European crust

Stefan Mroczek^{a,b,*}, Frederik Tilmann^{a,b}, Jan Pleuger^b, Xiaohui Yuan^a, Ben Heit^a, the SWATH-D Working Group¹, the AlpArray Working Group²

^a GFZ German Research Centre for Geosciences, Telegrafenberg, Potsdam, 14473, Brandenburg, Germany

^b Freie Universitaet Berlin, Institute for Geological Sciences, Malteserstr. 74-100, Berlin, 12249, Germany

ARTICLE INFO

Article history:

Received 28 September 2022

Received in revised form 22 February 2023

Accepted 27 February 2023

Available online 10 March 2023

Editor: H. Thybo

Keywords:

seismology

Moho map

receiver functions

Alps

ABSTRACT

The tectonic structure of the Eastern Alps is heavily debated with successive geophysical studies that are unable to resolve areas of ambiguity (e.g., the presence of a switch in subduction polarity and differing crustal models). In order to better understand this area, we produce a high resolution Moho map of the Eastern Alps based on a dense seismic broadband array deployment. Moho depths were derived from joint analysis of receiver function images of direct conversions and multiple reflections for both the SV (radial) and SH (transverse) components, which enables us to map overlapping and inclined discontinuities. We observe the European Moho to be underlying the Adriatic Moho from the west up to the eastern edge of the Tauern Window. East of the Tauern Window, a sharp transition from underthrusting European to a flat and thinned crust associated with Pannonian extension tectonics occurs, which is underthrust by both European crust in the north and by Adriatic crust in the south. The Adriatic lithosphere underthrusts northward below the Southern Alps and becomes steeper and deeper towards the Dinarides where it dips towards the north-east. Our results suggest that the steep high velocity region in the mantle below the Eastern Alps, observed in tomographic studies, is likely to be of European origin.

© 2023 Elsevier B.V. All rights reserved.

1. Introduction

Compared to oceanic subduction with a non-ambiguous sense of subduction and strongly localised deformation, continental convergence zones are characterized by wider deformation zones and interacting crustal blocks. This is especially true in the Eastern Alps where a stage of southward subduction of the Alpine Tethys was followed by collision of European and Adriatic continental lithosphere. During the final stages of orogeny, Eastern Alpine units escaped eastward into the opening Pannonian Basin (Ratschbacher et al., 1991) while at the same time the upper-plate Adriatic continental crust penetrated into the Alps.

Historically, the subduction of European lithosphere was assumed to be continuous across the entire Alps (e.g. Laubscher,

1971; Frisch, 1979), with the Adriatic microplate subducting in the east below the Dinarides (Laubscher, 1971). The subduction of European lithosphere has been confirmed with receiver function images of dipping Moho and Moho offsets in the Western and Central Alps (e.g. Lombardi et al., 2008; Spada et al., 2013; Zhao et al., 2015; Kummerow et al., 2004).

Teleseismic tomography results have consistently shown at least two main positive velocity anomalies in the upper mantle below the Alps (e.g. Babuška et al., 1990; Lippitsch et al., 2003; Mitterbauer et al., 2011; Karousová et al., 2013; Zhao et al., 2016; Paffrath et al., 2021; Plomerová et al., 2022). The western anomaly is accepted to be of European origin while the eastern anomaly has, in some studies, an apparent northerly dip leading to debate about its origin, either being of Adriatic or mixed European and Adriatic, rather than purely European, provenance (Babuška et al., 1990; Lippitsch et al., 2003; Karousová et al., 2013; Handy et al., 2015; Plomerová et al., 2022). This idea implies there is a change in subduction direction, i.e., a polarity switch from European subduction towards the south (in the Western and Central Alps) to Adriatic subduction towards the north in the Eastern Alps (linked to the Adriatic subduction beneath the Dinarides). Numerous studies probed the nature and location of this hypothetical polarity

* Corresponding author at: GFZ German Research Centre for Geosciences, Telegrafenberg, Potsdam, 14473, Brandenburg, Germany.

E-mail address: stefan.mroczek@outlook.com (S. Mroczek).

¹ Members of the SWATH-D Working Group who are not otherwise co-authors are M. Weber and C. Haberland.

² The complete member list for the AlpArray Working group can be found at alparray.ethz.ch.

switch and its geodynamic implications but have failed to provide conclusive results (e.g. Kissling et al., 2006; Ustaszewski et al., 2008; Handy et al., 2015). The TRANSALP profile (12°E), where a combination of passive (Kummerow et al., 2004) and controlled source (Lüschen et al., 2006) seismic imaging was employed, shows a gently southward dipping European Moho terminated by a jump to a shallow and sub-horizontal Adriatic Moho. The location of this transect is of particular interest because it falls almost exactly between the two main positive velocity anomalies described above.

The CELEBRATION 2000 (Guterch et al., 2003) and the ALP2002 (labelled ALP-01 and ALP-02 on Fig. 1) controlled source seismic (CSS) experiments reveal a region where thinned Pannonian crust (to the east) and the European Moho (showing a clear southward dip) intersect with Adriatic Moho to the south, suggesting that Europe remains the underlying plate (Brückl et al., 2010, 2007).

The best data coverage is provided along the N-S profile at 13.3°E by the passive EASI transect (AlpArray Seismic Network, 2014) which crosses a gap in the Moho map of Spada et al. (2013). The receiver function study carried out by Hetényi et al. (2018b), using the EASI temporary stations (Fig. 1) plus some permanent stations, showed the Adriatic and European Mohos forming a trough-like structure with a horizontal gap of nearly 50 km. They suggested that this gap does not contain a sharp Moho boundary but instead a ~20 km thick gradient zone reaching approximately 70 km depth. They connected this deeper gradient zone with the Adriatic Moho (making it consistent with Adriatic subduction) rather than with the European Moho.

An additional study probing the Moho and the mantle was conducted by Kind et al. (2021). Using stations that covered a much larger area, the method of stacking of S receiver functions without deconvolution was employed. This work not only mapped the Moho but also identified clear signals from negative velocity gradients in the mantle (NVGs), which have a dominant southward dip direction beneath the Alps. According to the authors, these interfaces (sometimes interpreted as the lithosphere-asthenosphere boundary) seem to be distributed in the entire Alpine region.

1.1. Outline of the tectonic evolution of the Eastern Alps and adjacent areas

The Eastern and eastern Southern Alps are a bivergent orogen consisting of a pro-wedge built up of units that were thrust northward onto the European foreland and a retro-wedge built up of units thrust southward onto Adria. From top to bottom, the pro-wedge consists of the Austroalpine nappes, the Penninic units derived from the Late Jurassic-mid-Cretaceous Alpine Tethys including small pieces of continental crust from within this oceanic basin, and Subpenninic and Helvetic units derived from the former European continental margin (e.g. Froitzheim et al., 2008; Schuster et al., 2013). Mostly in the Cretaceous, the Austroalpine units were assembled from the upper and lower plate of a probably intracontinental subduction zone within Adria (Stüwe and Schuster, 2010). In this eo-Alpine orogen, most of the sedimentary cover of the Austroalpine units was sheared off and piled up in an external position, forming the present-day Northern Calcareous Alps, while the basement units that were subducted deepest are now exposed in the axial parts of the Eastern Alps. The Alpine Tethys was subducted below Adria probably from the Early Cretaceous onwards (Faupl and Tollmann, 1979). Its suturing in the Eocene was followed by subduction of the thinned distal European continental margin and Oligocene collision of thicker European crust with Adria (e.g. Kurz et al., 2008). The occurrence of Penninic and Subpenninic units in the Eastern Alps is limited to a few tectonic windows. The largest of them, situated in the centre of our study area, is the Tauern Window (TW, Fig. 1) where Subpenninic units framed by Penninic ones are exposed below the

Austroalpine units. In the Neogene, European subduction in the Eastern Alps largely ceased as indicated by Neogene shortening amounts along the Northern Alpine Front of only a few kilometers (Hinsch, 2013). Instead, further convergence between the European and Adriatic plates was mostly accommodated by underthrusting of Adriatic crust below the pro-wedge, backthrusting in the Southern Alpine retro-wedge, and lateral extrusion of the Eastern Alps into the Pannonian Basin (Frisch et al., 1998). Major structures that accommodated this lateral extrusion are the dextral Periadriatic Fault along the boundary of the Eastern and Southern Alps and the sinistral Inntal and Salzachtal-Ennstal-Mariazell-Puchberg (SEMP) faults farther north as well as the Brenner and Katschberg normal faults at the western and eastern border of the TW, respectively (e.g. Linzer et al., 2002). The Austroalpine units continue eastwards into the internal and central Western Carpathians. In the external Western Carpathians, the narrow Pieniny Klippen Belt and the Magura Unit are lateral equivalents of the Penninic units (Froitzheim et al., 2008). The Magura Unit mostly consists of deep marine sediments that were deposited within the north-eastern part of Alpine Tethys (Magura Basin). Together with adjacent thin continental lithosphere of the distal European margin, the Magura Basin was part of a wide north-eastward-convex embayment in the European platform where flysch sedimentation continued into the Neogene (Kováč et al., 2017). While in the Alps relatively buoyant European continental crust had collided with Adria already in the Early Oligocene (Fig. 10a), slab-pull forces acting on dense lithosphere of the Magura Basin effected enhanced rollback of the Carpathian part of the European slab and fast retreat of the Carpathian trench towards the European foreland (Royden, 1993). This trench retreat allowed for the eastward extrusion (Ratschbacher et al., 1991) of mostly Austroalpine units in the AlCaPa (Alps-Carpathians-Pannonian) Mega-unit. This extrusion was concomitant with anticlockwise rotation of AlCaPa and with north-eastward advance and clockwise rotation of the Tisza Mega-unit into the southern Pannonian area. During extrusion into the Carpathian embayment, the AlCaPa and Tisza Mega-units were strongly stretched along their transport direction and thinned vertically (Ustaszewski et al., 2008). Diachronous collision of the AlCaPa and Tisza Mega-units with Europe is indicated by the ages of youngest thrusting in the external Carpathian flysch belt that range from ~18 Ma near the Alps-Carpathians junction to <10 Ma in the Eastern Carpathians (Jiříček, 1979). The Alpine tectonic evolution of the Dinarides started with Late Jurassic obduction of the West Vardar ophiolites onto the eastern Adriatic margin towards the Neotethys. After that, south-west-vergent thrusting of Adria-derived units was driven by north-eastward subduction of the Sava branch of Neotethys under Europe including Tisza. The north-western external Dinarides comprise, from top to bottom the Pre-Karst, High Karst, and Dalmatian units. These units are unmetamorphic to low-grade metamorphic and are characterised by Mesozoic sediments deposited on the Adriatic continental margin after the Middle Triassic onset of rifting that led to opening of the Neotethys. Most of the shortening in these units was accommodated during the Eocene to Oligocene, based on the age of syntectonic flysches exposed below thrust contacts. While the orogenic front shifted to the Dalmatian Unit in the Miocene, the hinterland was affected by Pannonian extension (Schmid et al., 2008). Neogene convergence between Adria and Europe in the area of the Alps-Dinarides junction was partitioned into south-vergent thrusts in the Southern Alps systematically overprinting older Dinaric structures (Doglioni, 1987), minor shortening along the Dinaric Frontal Thrust on Istria peninsula (~30 km since 15 Ma, van Hinsbergen et al., 2020), and a system of NW-SE striking dextral strike-slip faults between the Dinaric front and the Periadriatic Fault (Placer, 1998; Vrabec and Fodor, 2006). The kinematics of these faults is consistent with Neogene anticlockwise




Fig. 1. Simplified geological map of the study region with station locations plotted as triangles. Most stations belong to the dense SWATH-D and EASI deployments, which are supplemented by the temporary AlpArray Seismic Network (AASN) and permanent networks (with full citations in the supplemental material). Controlled source and/or passive seismic lines (NFP-20 East, TRANSALP, ALP-01, ALP-02, Alpass1-3, and AlpassDIP) are shown by grey lines. The Pannonian and associated basins are outlined in white in the inset map. AlCaPa - Alps-Carpathians-Pannonian Mega-unit, DFT - Dinaric Frontal Thrust, GC - Giudicarie Fault, IF - Idrija Fault, MU & PKB - Magura Unit & Pieniny Klippen Belt, NAF - Northern Alpine Front, PAF - Periadriatic Fault, SEMP - Salzach Ennstal Mariazell Puchberg fault, TW - Tauern Window, WVO - West Vardar Ophiolites. Tectonic map modified from M.R. Handy based on sources listed in Handy et al. (2019).

rotation of Adria with respect to Europe and increasing amounts of north-south convergence between the two plates towards the east. Ustaszewski et al. (2008) showed that a 20° anticlockwise rotation of Adria with respect to Europe around a pole near the north-western corner of Adria could potentially have placed a ~200 km long Adriatic slab under the Eastern Alps.

Von Blanckenburg and Davies (1995) proposed that after Europe-Adria collision, divergent forces between the shallower continental part of the European slab and the deeper and denser oceanic part led to breakoff of the latter at shallow depth, in turn triggering magmatism along the Periadriatic Fault. In terms of available radiometric ages (see compilations in Bergomi et al., 2015; Ji et al., 2019), Periadriatic magmatism peaked at ~35-30 Ma which, according to several authors (e.g. Schmid et al., 2013; Handy et al., 2015), is the time when the slab broke off. The change from underfilled (flysch-type) to overfilled (molasse-type) sedimentation in the Eastern Alpine foreland basin at ~20 Ma let Schlunegger and Kissling (2022) postulate that this slab breakoff happened only in the Early Miocene. According to these authors, Europe-Adria convergence in the Eastern Alps was accommodated by subduction of Adriatic lithosphere after ~20 Ma. Irrespective of its timing, breakoff of the European slab would be a necessary preliminary for a switch to Adriatic subduction under the Eastern Alps (e.g. Lippitsch et al., 2003; Kissling et al., 2006; Handy et al., 2015).

The migration of the Carpathian foreland depocentre has been interpreted to result from a subhorizontal tear propagating from west to east along which the European slab broke off (Nemcok et al., 1998; Wortel and Spakman, 2000), concentrating the slab pull force always on the westernmost undetached part of the slab and thereby causing subsidence in the foredeep. The eastward migration of the foreland depocentre started at ~19 Ma at the Alps-Carpathians junction and predated the cessation of thrusting (see above) always by ~1-2 Ma (Meulenkamp et al., 1996). Slab breakoff supposedly happened at shallow depth as evidenced by geophysical observations that the lithosphere below the Western Carpathians does not extend much farther down than the lithosphere of the European foreland (Bielik et al., 2022).

2. Data and methodology

In order to improve the data coverage in the Eastern Alps, the large SWATH-D deployment (Heit et al., 2021) was operated within the AlpArray initiative from summer 2017 to December 2019. Here, we analyse data provided by the SWATH-D experiment which consisted of 163 broadband stations in a wide E-W oriented network configuration with a nominal interstation distance of 15 km (Fig. 1). We exploited the dense coverage of the SWATH-D network as well as the concurrent AlpArray networks (Hetényi et al., 2018a) and the aforementioned EASI experiment to construct common conversion point (CCP) stacks (Kind et al., 2002) of receiver




Fig. 2. CCP stacks at 13.3°E (longitude of EASI) for the SV (a), SH (b, sensitive to dipping interfaces), and SV PpPs (c, migration for the first multiple) components. The lines show where the Moho interface has been picked. The solid line indicates that an interface was interpreted on that particular stack, while the dashed line indicates that it was picked from another. Traces from events with westerly back-azimuths have had their polarity flipped on the SH component. EU=European Moho and AD=Adriatic Moho.

functions (RFs), which helped to constrain the 3-D layout of the crust, including subducted fragments of lithosphere in the study region (Fig. 1).

We calculated RFs for the SV (S-wave vertical polarisation) and SH (S-wave horizontal polarisation) components from teleseismic P, PP, and PKP waveforms (see Kind and Yuan (2011) for a review and Sections S1 and S2 in the supplemental material for a more detailed description). The SV-RFs are standard tools for imaging horizontal interfaces, while SH-RFs are more sensitive to dipping structures. A subset of the used events and the back-azimuthal distribution for the full catalogue are plotted in Figure S1.

CCP stacks for both the SV and SH RFs were then assembled using a 1-D velocity model for the area (Jozi Najafabadi et al., 2021). In addition to CCP stacks for the primary conversion (P_s), we also calculated CCP stacks for the two main multiples (PpPs and PpSs+PsPs). For each cross section we show three CCP stacked images with the independent phases mentioned above (SV-P_s, SH-P_s, and SV-PpPs). As shown by synthetic tests (explained in detail

in Section 3), each of the three phases has different sensitivities to horizontal or dipping interfaces.

We finally identify the Moho interfaces below this complex tectonic region by combining all the CCP images with the guidance of the synthetic tests.

The Moho was picked manually, with the possibility of overlapping Moho conversions due to underthrust or subducted crustal sections. For convenience we make reference to the European, Adriatic, and Pannonian Mohos in order to fit appropriate overlapping interfaces. Identification of, in particular southward, dipping interfaces is challenging on the SV-P_s CCP stacks as expected conversion amplitudes are small (see Section 3), therefore we also relied on the SH-P_s and SV-PpPs stacks to identify those interfaces. To make a pick on the dipping sections of Moho, we required that coherent energy is observed on either the SV or SH component plus at least one of the multiples. For example, at 13.3°E and 14°E (Figs. 2 and 3) the energy on the SV component continues to depths greater than 150 km, however, we only pick to the extent observed on the first multiple. This way, for example, a diffraction




Fig. 3. CCP stack at 14°E as in Fig. 2. EU=European Moho, AD=Adriatic Moho, and PA=Pannonian Moho.

on the SV component extending the energy of a dipping interface beyond its true extent is less likely to be erroneously picked. These picks were then automatically re-picked and splines were fitted (with some manual adjustment) to obtain the Moho depth map. The spline coefficients, having been determined in the time domain, have units of Ps lag time and thus can be converted to splines for the first and second multiple with an appropriate Vp/Vs ratio. By maximising the fit of the spline with the multiple migrations, the average crustal Vp/Vs ratio was also determined. The normal CCP stacking procedure is based on a horizontally layered model, and will lead to artifacts when applied to phases converted at dipping interfaces (Cassidy, 1992), causing Vp/Vs to be overestimated and depth and dip to be underestimated. To account for this (to first order), we apply a dip-correction to the depths observed in the CCP stacks. For full details see Section S2. It should be noted that due to the selected event distribution, on the SH component, a positive velocity gradient (e.g. the Moho) will have positive polarity (red) for a south dipping interface but negative (blue) for a north dipping interface.

Indicative uncertainties in Moho depth are calculated by assuming a 0.3 s uncertainty in the Ps pick and a 25% uncertainty in

the dip angle, representing reasonable if somewhat arbitrary assumptions, which nevertheless illustrate how uncertainty in depth increases for inclined interfaces. Those error estimates do not include any errors due to lateral velocity variations.

3. Synthetic migration tests

The Moho geometry from different tectonic units beneath the study area is complex, with overlapping interfaces, horizontal, and strongly dipping segments present. As the event distribution is highly uneven (Figure S1), each of the RF phases/components has different sensitivities. Here we use synthetic RFs to evaluate the ability of different RF phases in recovering complicated Moho geometries. The target of the synthetic modelling is not to match the actual CCP stacks, but to provide clues about typical imaging artefacts and to test the capacity of our method to reconstruct the anomalies observed. In particular, we attempt to understand the role back-azimuthal distribution might play in the fidelity of dipping structures in the images obtained. We calculated synthetic migrations using the full wavefield finite difference method of Roecker et al. (2010) with a bandwidth of 1.25 s to 100 s. The de-




Fig. 4. Synthetic 2-layer crust and mantle models with realistic back-azimuthal distribution (assuming N-S orientation with dominant back-azimuth from the left/North). **A)** Synthetic trough model, note the dominance of the Moho dipping to the left. **B)** Based on the geometry observed at 14°E. Note the lack of coherent energy from right dipping interfaces. **C)** Same as A but for the first multiple. Note how the initial dipping portion of the Moho can be clearly seen as opposed to in B. Southward dipping phases at 450-600 km are the primary phase migrated upward.

convolution was performed with a fixed water level of 10%, and the CCP stacking was carried out assuming the IASP91 velocity model. V_p was set to 6.1 km/s for the crust and 8 km/s for the mantle with V_s set according to a constant $V_p/V_s=1.8$). Synthetics were calculated for a range of models with geometry simplified from the main interfaces we observe in the actual CCP images. The synthetics were computed in 2-D with oblique incidence angles (2.5-D). The synthetics represent the upper limit of resolvability for the Moho in the Alps for receiver functions with densely spaced stations (15 km inter-station distance), high signal-to-noise ratios, no intracrustal interfaces, and a two-layer (crust and upper mantle) velocity model. Thus the limitations in imaging stem largely from the irregular distribution of teleseismic events. To approximate the true event back-azimuth and slowness distribution while keeping the computational load moderate, the incidence angles and back-azimuths were clustered (detailed in the supplementary Section S3).

As mentioned, we consider the synthetics to represent the upper limit of our method to properly resolve the Moho. However, one aspect where the CCP stack of real data is likely to do bet-

ter than synthetics is with respect to diffractions. In the real Earth case, diffractions are expected to be much weaker, as the velocity variations and the interface geometries are not as sharp as in the synthetic models and also vary in 3-D rather than simply in 2-D (Lekić and Fischer, 2017). Thus, artefacts related to diffraction are likely exaggerated on the synthetic tests.

Comparing the results of the tests with our actual observations (Fig. 4), we observe some energy from Moho converted phases from the moderately dipping Moho on both the SV and SH components at 13.3°E (Fig. 2, north of 46.5°N). Further east, at 14°E (Fig. 3) we do not observe any energy from deep European Moho on the SV component below the flat section of the Pannonian Moho (south of 47.5°N). At first sight, this observation suggests the absence of subducted crust there. However, according to the synthetics shown in Fig. 4, no coherent energy would be expected on the SV component for this southward-directed dip angle. This is induced by the majority of event back-azimuths being from the N-E direction (Figure S1), so, for a more strongly southward dipping converter, incidence angles will be close to perpendicular and thus conversion coefficients will be very low. Therefore, we cannot




Fig. 5. CCP stack for 47.0°N with a swath width of 44 km. At approximately 13.4°E the jump from European to Pannonian Moho can be seen. Note that the marked European Moho beyond this longitude (from ~13.4°–14.2°) is not visible on the SV component CCP but instead was identified on the North-South SH and SV-PpPs CCP stacks. EU=European Moho and PA=Pannonian Moho.

conclude either for or against the presence of southward dipping European crust below the Pannonian Moho from the primary conversion (Ps phase) alone.

The event distribution should render a northerly dipping feature (such as the Adriatic Moho) easy to image due to high incidence angles resulting in large conversion coefficients such as at 14°E for the Adriatic Moho (Fig. 4). In contrast, a southward dipping European Moho would be very difficult to image (on the SV component) due to the almost complete lack of back-azimuths from the south. The situation is less ambiguous for multiples; for our northerly dominant back-azimuth distribution, the conversion coefficient for the multiple phases for a southward dipping converter would be higher than for the Ps phase (Fig. 4) and indeed we do observe a southward dipping European Moho for the first multiple (Fig. 3).

4. Results and discussion

Key CCP profiles are shown in Fig. 2 (N-S profile through Tauern window), Fig. 3 (N-S profile just east of the Tauern window) and Fig. 5 (E-W profile). Figs. 6 and 7 compare the N-S profiles with relevant synthetics and highlight the inferred Moho geometry corrected for dip. A larger suite of profiles can be found in the supplementary Section S6, Figures S10–S16. The Moho map is plotted in Fig. 8. All three domains have on average a more felsic composition with average V_p/V_s ratios of 1.72 ± 0.03 for Adriatic, 1.70 ± 0.04 European, and 1.70 ± 0.03 for Pannonian crust. As V_p/V_s values at specific locations are sensitive to the dip correction and thus sometimes poorly constrained, we do not plot the full V_p/V_s ratio map to avoid misinterpretation. As demonstrated by the synthetic migrations (Section 3), we expect interfaces dipping towards the south to be difficult to image while those dipping towards the north will be more dominant.

4.1. Moho geometry

In the west of the study area (up to 13°E), our results do not differ significantly from previous studies. The NFP-20 East CSS survey (Valasek et al., 1991, see Fig. 1 for location) shows similar results to the section at 10°E (Figure S10). At 12°E, our results agree well with those provided by the TRANSALP experiment (Kummerow et al., 2004) (Figure S12).

Our synthetic modelling (Fig. 4) demonstrates that, given a realistic teleseismic event distribution, a simple Moho trough model with no sense of subduction could give rise to an apparent pref-

erence for northward subduction due to the stronger conversions for a north dipping interface, illuminated from predominantly northerly azimuths (Fig. 6). For the European Moho alone there is little disagreement with Hetényi et al. (2018b) down to ~60 km depth. We go on to interpret deeper energy to be from the European Moho, based on the SH component and multiples. In contrast, in our interpretation the Adriatic Moho terminates much earlier, at approximately 40 km depth, similar to the Moho geometry of Spada et al. (2013). This disagreement in interpretation stems largely from our findings of a southward dipping interface on migrations for the SH component and multiples aided by a larger suite of stations/events and a wider E-W extent.

With higher velocity mantle emplaced over subducted crust, one would expect a negative velocity contrast (blue band), representing the top of the crustal sliver, to be present above the positive contrast deeper Moho interface (red band), including on the SH component (e.g. Abe et al., 2011). At 13.3°E (Fig. 6), a coherent south dipping negative interface can be seen above the dipping interval of the European Moho on the SH component which may correspond to the mantle to subducted crust transition. On the migration for the first multiple, a negative polarity signal with similar amplitude to the PpPs phase can also be observed. At 14°E (Fig. 7), this interface is clear on the migration for the first multiple but appears disturbed on the SH component.

4.2. Transition to the Pannonian region

The transect at 14.0°E (Fig. 7) shows a distinctly different crustal structure from those to the west. A flat Moho at ~35 km depth between 47.5°N and 46.5°N is underthrust by the European Moho from the north and by the Adriatic Moho from the south. This section of crust has presumably been thinned and reshaped during lateral extrusion into the Pannonian Basin (Ratschbacher et al., 1991), with possible additional removal of crust by delamination, which would imply the interface represents a new Moho. For convenience, we refer to this interface as Pannonian Moho, although its original provenance is Adriatic.

This thinned crust has been observed previously in the Eastern Alps by Brückl et al. (2010). With the CSS data from the ALP2002 experiment, Brückl et al. (2007) suggested a westward termination of the Pannonian Moho at approximately 14°E and then further refined this to ~13.6°E with 2-D elastic plate modelling (Brückl et al., 2010), approximately at the longitude of the eastern edge of the Tauern Window. We find the westward termination of the flat Moho section to be close to the point inferred by Brückl et al.




Fig. 6. SV CCP stacks for 13.3°E with a swath width of 44 km and with the SH component (the polarities of events with back-azimuths from the west are flipped) inset and the first multiple (PpPs, below). Both the SH component and the multiple are more sensitive to dipping layers which are not observed on the SV component (Fig. 2 shows the unaltered images). Red indicates a velocity increase downwards (e.g. the Moho) and blue a velocity decrease. The SV synthetic illustrates the difficulty of imaging southward dipping interfaces, making northward dipping interfaces more dominant. EU=European Moho and AD=Adriatic Moho.




Fig. 7. As Fig. 6 for the CCP stack at 14°E (unaltered Fig. 3). The SV synthetic illustrates why the lack of energy on the true SV component is expected due to a combination of steeply dipping interfaces and lack of illumination from the south. EU=European Moho, AD=Adriatic Moho, and PA=Pannonian Moho.




Fig. 8. Moho depth map (A) showing the three main Moho domains. Paler areas show interpolated depths (no manual picks). The color coded map refers to the shallowest Moho in cases of overlap. Additionally, 10 km contour lines are shown for each domain that highlight the deeper Moho in case of overlaps due to underthrusting. Moving from west to east, the European Moho (purple) underthrusts the shallowly dipping Adriatic Moho (blue). At $\sim 13^\circ\text{E}$, Adriatic Moho begins to retreat southwards into the Dinarides as it underthrusts extended Pannonian crust. Likewise, the European Moho dips steeply beneath the Pannonian Moho (green) to a depth of ~ 150 km. **B**) shows the same Moho geometry in 3-D from north-easterly aspect for the same extent as **A**). Dipping regions of the European Moho where we do not observe energy on either component or the multiples are masked with black.

(2010) (approximately 13.5°E visible on the E-W profile at 47.0°N -Fig. 5), which is imaged directly in the CSS data of the E-W ALP '75 profile (Yan and Mechie, 1989). The interpretation of this interface as being the Moho is confirmed by the ambient noise surface wave tomography of Lu et al. (2020), where it coincides with the 4.2 km/s Moho contour (shown by a dotted line on Fig. 7), implying typical mantle velocities below. The western most extent of the thinned crust also coincides with a sharp increase in gravity moving eastwards out of the Tauern Window (Zahorec et al., 2021).

The visually step-like structure between the European and Pannonian Mohos coincides spatially with the SEMP fault, which marks the northerly extent of the lateral eastward extrusion of the Eastern Alps into the Pannonian Basin (Ratschbacher et al., 1991). A similar observation, using CSS data, was made further east across the Carpathian Arc where the dipping Moho interface from the European platform intersects the flat Moho below the Pannonian Basin (Tomek and Hall, 1993; Bielik et al., 2004).

Below the European to Pannonian Moho step, we image a sharply dipping interface that can be seen clearly on the SH component and the SV multiples. We interpret this interface as being continuous with the European Moho at least until 13.5°E (moving eastwards) and the onset of the Pannonian Moho. There is a small gap at 60 km depth between that and the flat European Moho we observe on the SV component at 14°E . This gap is perhaps real or may be an artefact produced by a rather low conversion coefficient (the conversion coefficient depends on the angle of incidence). We choose to draw a connected interface based on continuity with migrations to the west but distinguish this as a region of high uncertainty.

Ongoing Adriatic subduction, with little shortening after the Oligocene, forms the Dinarides (Sun et al., 2019) to the south of our study area. As we move east from the flat Adriatic Moho we observe in the west, this transition must be reflected in the Moho geometry. By 14°E , the Adriatic Moho retreats and begins to dip below the Pannonian Moho. We observe a step up from the Adriatic Moho to the Pannonian Moho at 46.25°N with approximately 20 km offset in depth (Fig. 7). The Adriatic Moho may also underthrust the Pannonian Moho but only on the order of a few tens kilometers. We have high confidence in this observa-

tion as northerly back-azimuths give high conversion coefficients (as demonstrated in the synthetic tests in Fig. 4).

4.3. Comparison with teleseismic tomography

In Figures S17-S19, we compare the tomographic model of Paf-rath et al. (2021) (from teleseismic P-wave tomography) to our observations along our main transects. Handy et al. (2021) interpret the European slab in this model to be detached, contrary to the interface we observe with receiver functions. The simultaneous teleseismic tomography study of Plomerová et al. (2022) instead interpret this anomaly to be undetached and of Adriatic or mixed Adriatic/European origin. Potential reasons for this difference are discussed by Plomerová et al. (2022). Additionally, receiver functions image velocity contrasts so a layer well below the resolution limit of the tomography could still produce a clear conversion. In the light of disagreement between concurrent tomographic models and differences in imaging capabilities, we believe that an interface connecting the anomaly at depth with European crust cannot be ruled out based on teleseismic travel time tomography alone. Full waveform inversion or joint inversion may be required to resolve these differences.

We plot the observations of both tomographic studies and compare them to our Moho interface in Fig. 9. The NVG, observed by Kind et al. (2021) at 14°E , almost exactly parallels the Moho interface we trace (before depth correction) reaching the central positive velocity anomaly observed in both studies.

5. Tectonic implications

West of 13.5°E , the Moho geometries described before (Section 4.1) are well in line with the classical tectonic concept of the Alps consisting of a pro-wedge and a retro-wedge underlain by European and Adriatic Mohos, respectively. Farther east, the Moho configuration imaged in Fig. 3 requires that lower crust is still attached to the downgoing European and Adriatic plates while the former upper crust, sheared off from these plates, overlies the Pannonian Moho (Fig. 10c). Since the European and Adriatic Mohos dip below the Pannonian one, the undetached lower crust of




Fig. 9. P-wave tomography at 14°E (Paffrath et al., 2021) (A) and at 14.2°E (Plomerová et al., 2022) (B). The solid black outline shows where the sign of the anomaly is opposite between the two models. This highlights the central positive slab anomaly which has no clear preferred dip direction. The NVG (lower dot-dashed line) of Kind et al. (2021) (uncorrected for dip/depth) and upper receiver function boundary lead (solid green) into where both studies observe a positive anomaly at depth. EU=European Moho, AD=Adriatic Moho, and PA=Pannonian Moho.

both plates must be separated from the Pannonian Moho by intervening mantle material. We hypothesise that emplacement of the mantle material was the combined effect of a retreat of the Alpine segment of the European slab with respect to the Adriatic upper plate and rollback of the Carpathian segment of the European slab that caused strong thinning of the Alpine crust in the upper plate. Retreat of the European slab below the Alps may have already started when the Alpine Tethys was sutured or soon thereafter (Fig. 10a), similar to scenarios proposed by Malusà et al. (2011) and Kissling and Schlunegger (2018) for the Western and Central Alps. In accordance with classical models of Alpine nappe formation (e.g. Schmid et al., 1996), we assume that the Subpenninic nappes were successively stacked upon each other along in-sequence thrusts but that, different from classical models, mantle material migrated along the décollement between the rear parts of the Subpenninic nappes and the subducting, non-detached European lower crust. The emplacement of mantle material below Penninic and Subpenninic nappes of the Alpine pro-wedge probably started when the Alpine Tethys and lithosphere, comprising only thin crust of the former distal European passive margin, were subducted. At this stage (~ 32 Ma, Fig. 10a), asthenosphere upwelling above the European slab would be a potential cause of magmatism along the Periadriatic Fault (Ji et al., 2019). We assume that the northward advance of the Pannonian Moho was concomitant with steepening and delamination of the European slab because after suturing of the Alpine Tethys in the Eocene, the Northern Alpine Front hardly moved towards the foreland and became almost inactive in the Neogene. At 14°E , the length of European Moho underthrust below the Alps far exceeds estimates of Neogene thrusting along the Northern Alpine Front and so must have been subducted mostly in pre-Neogene times. Together with steepening of the European slab, we consider eastward escape of Eastern Alpine orogenic crust into the Pannonian basin from the Miocene on as the main driver behind the formation of the Pannonian Moho. Considering values of 28%–39% and 47% for Miocene N-S directed shortening and E-W directed extension, respectively, in the Eastern Alps (Linzer et al., 2002), the eastward escape thinned the crust in the axial part of the Alps more efficiently than north-south convergence between Europe and Adria could have thickened it. We note that according to several au-

thors, the Adriatic lower crust had started to underthrust below the evolving Southern Alps, causing N-S shortening also in the Eastern Alps in the Late Oligocene–Early Miocene (Schmid et al., 2013), before the Alps were affected by Carpathian rollback since the Early to Middle Miocene (Schmid et al., 2013). Irrespective of the relative timing of Adriatic underthrusting and Carpathian rollback, the presence of a flat Moho from below the eastern Tauern Window connecting seamlessly eastwards into the Moho below the Pannonian basin supports the notion that the flat Pannonian Moho was mostly shaped in conjunction with Carpathian rollback. We consider our finding that the European Moho below the Eastern Alps extends below the northern edge of the Adriatic Moho as evidence that no subduction polarity switch occurred below the Eastern Alps. Given the fact that the NVG of Kind et al. (2021) connects to the upper boundary of the positive P-wave velocity anomaly, as imaged by Paffrath et al. (2021) and Plomerová et al. (2022), and that the European Moho can be traced downward into the same anomaly (Fig. 9), the European slab appears to be continuous to at least the bottom of the positive P-wave anomalies, i.e. a depth 200 km. This implies that the Alpine part of the European slab, contrary to the Carpathian part, did not break off at shallow depth in the Neogene. We therefore suggest that the change from underfilled to overfilled conditions in the foreland basin of the Eastern Alps was not the response to shallow slab breakoff in the Early Miocene (Schlunegger and Kissling, 2022), but resulted rather from a decreasing topographic load exerted by the Alpine orogenic crust. It is feasible that this decrease was caused by the eastward escape of Alpine orogenic crust into the Pannonian basin, thereby reducing the area occupied by this crust in cross section (compare Figs. 10b and 10c). If this is correct, 20 Ma is the latest possible onset for lateral escape of AlCaPa driven by rollback of the Carpathian part of the European slab.

The European Moho extending below the northern edge of the Adriatic Moho limits the possible amount of Neogene Adriatic underthrusting below the Alps to a few tens of kilometres. According to our Moho map (Fig. 8), the underthrust Adriatic Moho below the Eastern Alps directly continues to below the Dinarides where it underlies the Pannonian Moho. The Adriatic Moho can be traced until ~ 75 km north-east of the Dinaric Frontal Thrust (Fig. 8a) and until ~ 30 km north-east of the south-western edge of the




Fig. 10. Schematic cross sections through the Alps at 14°E illustrating the tectonic evolution of the Pannonian Moho. The restorations are not balanced because of large out-of-plane movements of tectonic units. **A)** Onset of steepening of the European slab after subduction of the Alpine Tethys and distal European margin. Note the presence of a detachment and imbricated structures within the European plate, which will be added to the European accreted units. **B)** Continued steepening of the European slab and onset of eastward escape of the Eastern Alps between the SEMP and Periadriatic Fault led to northward propagation of the Pannonian Moho by insertion of mantle material between the Subpenninic units and undetached European lower crust. Though compared to their present-day geometry, the Eastern Alps were shortened by ~65 km mostly in the Northern Calcareous Alps and just north of the Periadriatic Fault (Linzer et al., 2002), the crust above the Pannonian Moho was not thickened significantly due to the effect of eastward escape. The scissor icon indicates decoupling. **C)** Present situation, Moho configuration (purple lines) is taken from Fig. 7. To the south, the Pannonian Moho extends below the active Dinaric strike-slip fault system of which only the Idrija fault is shown here. DFT – Dinaric Frontal Thrust, IF – Idrija Fault, NAF – Northern Alpine Front, NCA – Northern Calcareous Alps, PAF – Periadriatic Fault, SAT – South Alpine frontal thrust, SEMP – Salzach-Ennstal-Mariazell-Puchberg Fault, SF – Sava Fault.

Pannonian Moho. Comparing this to the post-15 Ma north-east to south-west shortening of ~30 km by van Hinsbergen et al. (2020), more than half of the Adriatic underthrusting should have occurred before 15 Ma. Assuming that the Adriatic Moho does not extend farther to the north-east than imaged in Fig. 8 and that the shortening estimate of ~30 km by van Hinsbergen et al. (2020) is correct, underthrusting of the Adriatic under the Pannonian Moho would have happened during the last 15 Ma. This would comply with continued Miocene underthrusting at the Dinaric Frontal Thrust while, in the hinterland, the Pannonian Moho was formed by extension since the Early Miocene. In Fig. 10, we assume that Late Oligocene-Early Miocene thrusting in the Southern Alps led to thickening of the Adriatic crust between the Dinaric Frontal Thrust and the Periadriatic Fault and that this crust was thinned again by transtensional faulting in the same area (Vrabec and Fodor, 2006). A system of dextral strike-slip faults in the external Dinarides, that run roughly parallel the Dinaric Frontal Thrust and that are partly still active today, coincides with the south-western limit of the Pannonian Moho below the Dinarides.

6. Conclusions

The SWATH-D and AlpArray seismic networks have allowed the Moho in the Eastern Alps to be delineated to an unprecedented degree of resolution. We are able to shed light on crustal configuration and produce a map of the Moho depth of this region, in some places distinguishing overlapping Mohos. The Moho map, determined from receiver functions, shows a thinned Pannonian (originally Adriatic) crust east of the Tauern Window. At this longitude (14°E), a step from the Adriatic Moho to the Pannonian Moho suggests underthrusting, but only on the order of a few kilometers, while the European Moho dips steeply beneath the Pannonian Moho down to a depth of at least 150 km depth. Moving west, the deeper subducted European crust shallows out to ~70 km depth by 12°E (TRANSALP), remaining the lower plate but now underthrusting directly beneath the shallowly dipping Adriatic Moho.

This configuration of the European, Adriatic, and Pannonian Mohos is incompatible with a switch of subduction polarity in the Eastern Alps. The Pannonian Moho formed due to enhanced roll-

back subduction of the European slab below the Carpathians and back-arc extension in the Pannonian Basin that caused lateral escape of the Eastern Alps towards the Pannonian Basin. Steepening and delamination of the European slab along the top of undetached, subducting lower crust was a preliminary for the expansion of the Pannonian Moho into the Eastern Alps that probably started no later than 20 Ma. The Pannonian Moho also extends well into the External Dinarides. Limited amounts of underthrusting of the Adriatic Moho under the Pannonian Moho are in line with Miocene shortening amounts estimated for the north-western Dinarides.

7. Data and resources

Seismic data used for this publication are hosted at the GEOFON data centre (SWATH-D) and other data centres within EIDA (AlpArray data and permanent stations), see Section S5 for a full list of data references and network codes. The Moho map, equivalent Ps lag time map for the Moho conversion, and map of implied Vp/Vs are made available at doi:<https://doi.org/10.5880/GFZ.2.4.2021.009>. The hill shade map was calculated from the Copernicus EU-DEM (produced with funding by the European Union, doi:<https://doi.org/10.5270/ESA-c5d3d65>).

CRediT authorship contribution statement

Stefan Mroczek: Formal analysis, Methodology, Writing – original draft. **Frederik Tilmann:** Supervision, Writing – review & editing. **Jan Pleuger:** Writing – review & editing. **Xiaohui Yuan:** Writing – review & editing. **Ben Heit:** Writing – review & editing.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data will be made available on request.

Acknowledgements

We thank Mark Handy, György Hetényi, Rainer Kind, and Stefan Schmid for in depth discussions and valuable feedback as well as the AlpArray-EASI Working Group for their work on the EASI experiment and the AlpArray Working Group on the AlpArray Seismic Network. We would also like to extend our thanks to the Deutsche Forschungsgemeinschaft [Grant numbers TI 316/8-1, TI 316/601, YU 115/11-1, and PL 534/4-1] for the funding of this research. We thank Steven Roecker for making available his software WT_MPI used for synthetics calculation (Roecker et al., 2010).

We acknowledge and thank the SWATH-D Field Team (the complete list can be found at doi:<https://doi.org/10.14470/MF7562601148>) for the data collection). We also thank Tuna Eken, one anonymous reviewer, and associate editor Hans Thybo for their constructive comments which helped to improve the clarity of presentation.

Appendix A. Supplementary material

Supplementary material related to this article can be found online at <https://doi.org/10.1016/j.epsl.2023.118096>.

References

Abe, Y., Ohkura, T., Hirahara, K., Shibutani, T., 2011. Common-conversion-point stacking of receiver functions for estimating the geometry of dipping interfaces. *Geophys. J. Int.* 185, 1305–1311. <https://doi.org/10.1111/j.1365-246X.2011.05001.x>.

- AlpArray Seismic Network, 2014. Eastern Alpine seismic investigation (EASI) – AlpArray complimentary experiment. https://doi.org/10.12686/ALPARRAY/XT_2014.
- Babuška, V., Plomerova, J., Granet, M., 1990. The deep lithosphere in the Alps: a model inferred from P residuals. *Tectonophysics* 176, 137–165. [https://doi.org/10.1016/0040-1951\(90\)90263-8](https://doi.org/10.1016/0040-1951(90)90263-8).
- Bergomi, M.A., Zanchetta, S., Tunesi, A., 2015. The Tertiary dike magmatism in the Southern Alps: geochronological data and geodynamic significance. *Int. J. Earth Sci.* 104, 449–473. <https://doi.org/10.1007/s00531-014-1087-5>.
- Bielik, M., Šefara, J., Kováč, M., Bezák, V., Plašienka, D., 2004. The Western Carpathians–interaction of Hercynian and Alpine processes. *Tectonophysics* 393, 63–86. <https://doi.org/10.1016/j.tecto.2004.07.044>.
- Bielik, M., Zeyen, H., Starostenko, V., Makarenko, I., Legostaeva, O., Savchenko, S., Dérerová, J., Grinč, M., Godová, D., Pánisová, J., 2022. A review of geophysical studies of the lithosphere in the Carpathian-Pannonian region. *Geol. Carpath.* 73. <https://doi.org/10.31577/GeolCarp.73.6.2>.
- von Blanckenburg, F., Davies, J.H., 1995. Slab breakoff: a model for syncollisional magmatism and tectonics in the Alps. *Tectonics* 14, 120–131. <https://doi.org/10.1029/94TC02051>.
- Brückl, E., Behm, M., Decker, K., Grad, M., Guterch, A., Keller, G., Thybo, H., 2010. Crustal structure and active tectonics in the Eastern Alps. *Tectonics* 29. <https://doi.org/10.1029/2009TC002491>.
- Brückl, E., Bleibinhaus, F., Gosar, A., Grad, M., Guterch, A., Hrubcová, P., Keller, G.R., Majdaňski, M., Šumanovac, F., Tiira, T., et al., 2007. Crustal structure due to collisional and escape tectonics in the Eastern Alps region based on profiles Alp01 and Alp02 from the ALP 2002 seismic experiment. *J. Geophys. Res.* 112. <https://doi.org/10.1029/2006JB004687>.
- Cassidy, J., 1992. Numerical experiments in broadband receiver function analysis. *Bull. Seismol. Soc. Am.* 82, 1453–1474. <https://doi.org/10.1785/BSSA0820031453>.
- Dogliioni, C., 1987. Tectonics of the Dolomites (Southern Alps, Northern Italy). *J. Struct. Geol.* 9, 181–193. [https://doi.org/10.1016/0191-8141\(87\)90024-1](https://doi.org/10.1016/0191-8141(87)90024-1).
- Faupl, P., Tollmann, A., 1979. Die Roßfeldschichten: Ein Beispiel für Sedimentation im Bereich einer tektonisch aktiven Tiefseerinne aus der kalkalpinen Unterkreide. *Geol. Rundsch.* 68, 93–120. <https://doi.org/10.1007/BF01821124>.
- Frisch, W., 1979. Tectonic progradation and plate tectonic evolution of the Alps. *Tectonophysics* 60, 121–139. [https://doi.org/10.1016/0040-1951\(79\)90155-0](https://doi.org/10.1016/0040-1951(79)90155-0).
- Frisch, W., Kuhlemann, J., Dunkl, I., Brügel, A., 1998. Palinspastic reconstruction and topographic evolution of the Eastern Alps during late Tertiary tectonic extrusion. *Tectonophysics* 297, 1–15. [https://doi.org/10.1016/S0040-1951\(98\)00160-7](https://doi.org/10.1016/S0040-1951(98)00160-7).
- Froitzheim, N., Plašienka, D., Schuster, R., 2008. Alpine tectonics of the Alps and Western Carpathians. In: *The Geology of Central Europe Volume 2: Mesozoic and Cenozoic*. Geological Society of London.
- Guterch, A., Grad, M., Keller, G., Posgay, K., Vožar, J., Špičák, A., Brückl, E., Hajnal, Z., Thybo, H., Selvi, O., et al., 2003. Celebration 2000 seismic experiment. *Stud. Geophys. Geod.* 47, 659–669.
- Handy, M.R., Giese, J., Schmid, S.M., Pleuger, J., Spakman, W., Onuzi, K., Ustaszewski, K., 2019. Coupled crust-mantle response to slab tearing, bending, and rollback along the Dinaride-Hellenide orogen. *Tectonics* 38, 2803–2828. <https://doi.org/10.1029/2019TC005524>.
- Handy, M.R., Schmid, S.M., Paffrath, M., Friederich, W., the AlpArray Working Group, 2021. Orogenic lithosphere and slabs in the greater Alpine area – interpretations based on teleseismic P-wave tomography. *Solid Earth* 12, 2633–2669. <https://doi.org/10.5194/se-12-2633-2021>.
- Handy, M.R., Ustaszewski, K., Kissling, E., 2015. Reconstructing the Alps–Carpathians–Dinarides as a key to understanding switches in subduction polarity, slab gaps and surface motion. *Int. J. Earth Sci.* 104, 1–26. <https://doi.org/10.1007/s00531-014-1060-3>.
- Heit, B., Cristiano, L., Haberland, C., Tilmann, F., Pesaresi, D., Jia, Y., Hausmann, H., Hemmleb, S., Haxter, M., Zieke, T., Jaeckl, K., Schloemer, A., Weber, M., 2021. The SWATH-D seismological network in the Eastern Alps. *Seismol. Res. Lett.* 92, 1592–1609. <https://doi.org/10.1785/0220200377>.
- Hetényi, G., Molinari, I., Clinton, J., Bokelmann, G., Bondár, I., Crawford, W.C., Dessa, J.X., Doubre, C., Friederich, W., Fuchs, F., et al., 2018a. The AlpArray seismic network: a large-scale European experiment to image the Alpine orogen. *Surv. Geophys.* <https://doi.org/10.1007/s10712-018-9472-4>.
- Hetényi, G., Plomerová, J., Bianchi, I., Exnerová, H.K., Bokelmann, G., Handy, M.R., Babuška, V., AlpArray-EASI Working Group and others, 2018b. From mountain summits to roots: crustal structure of the Eastern Alps and Bohemian Massif along longitude 13.3°E. *Tectonophysics* 744, 239–255. <https://doi.org/10.1016/j.tecto.2018.07.001>.
- van Hinsbergen, D.J.J., Torsvik, T.H., Schmid, S.M., Mačenco, L.C., Maffione, M., Vissers, R.L.M., Güler, D., Spakman, W., 2020. Orogenic architecture of the Mediterranean region and kinematic reconstruction of its tectonic evolution since the Triassic. *Gondwana Res.* 81, 79–229. <https://doi.org/10.1016/j.gr.2019.07.009>.
- Hinsch, R., 2013. Laterally varying structure and kinematics of the Molasse fold and thrust belt of the Central Eastern Alps: implications for exploration. *Am. Assoc. Pet. Geol. Bull.* 97, 1805–1831. <https://doi.org/10.1306/04081312129>.
- Ji, W.Q., Malusà, M.G., Tiepolo, M., Langone, A., Zhao, L., Wu, F.Y., 2019. Synchronous periadriatic magmatism in the Western and Central Alps in the absence of slab breakoff. *Terra Nova* 31, 120–128. <https://doi.org/10.1111/ter.12377>.

- Jiříček, R., 1979. Tectonic development of the Carpathian arc in the Oligocene and Neogene. In: *Tectonic Profiles Through the Western Carpathians*. Geological Institute Dionyz Stur, Bratislava, pp. 205–214.
- Jozsi Najafabadi, A., Haberland, C., Ryberg, T., Verwater, V.F., Le Breton, E., Handy, M.R., Weber, M., 2021. the AlpArray and AlpArray SWATH-D working groups. Relocation of earthquakes in the southern and eastern Alps (Austria, Italy) recorded by the dense, temporary SWATH-D network using a Markov chain Monte Carlo inversion. *Solid Earth* 12, 1087–1109. <https://doi.org/10.5194/se-12-1087-2021>.
- Karousová, H., Plomerová, J., Babuška, V., 2013. Upper-mantle structure beneath the southern Bohemian Massif and its surroundings imaged by high-resolution tomography. *Geophys. J. Int.* 194, 1203–1215. <https://doi.org/10.1093/gji/ggt159>.
- Kind, R., Schmid, S.M., Yuan, X., Heit, B., Meier, T., the AlpArray and AlpArray-SWATH-D Working Groups, 2021. Moho and uppermost mantle structure in the Alpine area from S-to-P converted waves. *Solid Earth* 12, 2503–2521. <https://doi.org/10.5194/se-12-2503-2021>.
- Kind, R., Yuan, X., 2011. Seismic receiver function technique. In: Gupta, H. (Ed.), *Encyclopedia of Solid Earth Geophysics*. Springer, pp. 1258–1269.
- Kind, R., Yuan, X., Saul, J., Nelson, D., Sobolev, S., Mechie, J., Zhao, W., Kosarev, G., Ni, J., Achauer, U., et al., 2002. Seismic images of crust and upper mantle beneath Tibet: evidence for Eurasian Plate subduction. *Science* 298, 1219–1221. <https://doi.org/10.1126/science.1078115>.
- Kissling, E., Schlunegger, F., 2018. Rollback orogeny model for the evolution of the Swiss Alps. *Tectonics* 37, 1097–1115. <https://doi.org/10.1002/2017TC004762>.
- Kissling, E., Schmid, S.M., Lippitsch, R., Ansoerge, J., Fügenschuh, B., 2006. Lithosphere structure and tectonic evolution of the Alpine arc: new evidence from high-resolution teleseismic tomography. *Geol. Soc. Lond. Mem.* 32, 129–145. <https://doi.org/10.1144/GSL.MEM.2006.032.01.08>.
- Kováč, M., Márton, E., Oszczytko, N., Vojtko, R., Hók, J., Králiková, S., Plašienka, D., Klučiar, T., Hudáčiková, N., Oszczytko-Clowes, M., 2017. Neogene palaeogeography and basin evolution of the Western Carpathians, Northern Pannonian domain and adjoining areas. *Glob. Planet. Change* 155, 133–154. <https://doi.org/10.1016/j.gloplacha.2017.07.004>.
- Kummerow, J., Kind, R., Oncken, O., Giese, P., Ryberg, T., Wylegalla, K., Scherbaum, F., Group, T.W., et al., 2004. A natural and controlled source seismic profile through the Eastern Alps: TRANSALP. *Earth Planet. Sci. Lett.* 225, 115–129. <https://doi.org/10.1016/j.epsl.2004.05.040>.
- Kurz, W., Handler, R., Bertoldi, C., 2008. Tracing the exhumation of the eclogite zone (tauern window, eastern alps) by ⁴⁰Ar/³⁹Ar dating of white mica in eclogites. In: *Orogenic Processes in the Alpine Collision Zone*. Springer, pp. S191–S206.
- Laubscher, H., 1971. Das Alpen-Dinariden-Problem und die Palinspastik der südlichen Tethys. *Geol. Rundsch.* 60, 813–833. <https://doi.org/10.1007/BF02046522>.
- Lekić, V., Fischer, K.M., 2017. Interpreting spatially stacked Sp receiver functions. *Geophys. J. Int.* 210, 874–886. <https://doi.org/10.1093/gji/ggx206>.
- Linzer, H.G., Decker, K., Peresson, H., Dell'Mour, R., Frisch, W., 2002. Balancing lateral orogenic float of the eastern alps. *Tectonophysics* 354, 211–237. [https://doi.org/10.1016/S0040-1951\(02\)00337-2](https://doi.org/10.1016/S0040-1951(02)00337-2).
- Lippitsch, R., Kissling, E., Ansoerge, J., 2003. Upper mantle structure beneath the Alpine orogen from high-resolution teleseismic tomography. *J. Geophys. Res.* 108. <https://doi.org/10.1029/2002JB002016>.
- Lombardi, D., Braunmiller, J., Kissling, E., Giardini, D., 2008. Moho depth and Poisson's ratio in the Western-Central Alps from receiver functions. *Geophys. J. Int.* 173, 249–264. <https://doi.org/10.1111/j.1365-246X.2007.03706.x>.
- Lu, Y., Stehly, L., Brossier, R., Paul, A., AlpArray Working Group, 2020. Imaging Alpine crust using ambient noise wave-equation tomography. *Geophys. J. Int.* 222, 69–85. <https://doi.org/10.1093/gji/ggaa145>.
- Lüschen, E., Borrini, D., Gebrande, H., Lammerer, B., Millahn, K., Neubauer, F., Nicolich, R., Group, T.W., et al., 2006. TRANSALP—deep crustal Vibroseis and explosive seismic profiling in the Eastern Alps. *Tectonophysics* 414, 9–38. <https://doi.org/10.1016/j.tecto.2005.10.014>.
- Malusà, M.G., Faccenna, C., Garzanti, E., Polino, R., 2011. Divergence in subduction zones and exhumation of high pressure rocks (Eocene Western Alps). *Earth Planet. Sci. Lett.* 310, 21–32. <https://doi.org/10.1016/j.epsl.2011.08.002>.
- Meulenkamp, J., Kováč, M., Cicha, I., 1996. On Late Oligocene to Pliocene depocentre migrations and the evolution of the Carpathian-Pannonian system. *Tectonophysics* 266, 301–317. [https://doi.org/10.1016/S0040-1951\(96\)00195-3](https://doi.org/10.1016/S0040-1951(96)00195-3).
- Mitterbauer, U., Behm, M., Brückl, E., Lippitsch, R., Guterch, A., Keller, G.R., Koslovskaya, E., Rumpfhuber, E.M., Šumanovac, F., 2011. Shape and origin of the East-Alpine slab constrained by the ALPASS teleseismic model. *Tectonophysics* 510, 195–206. <https://doi.org/10.1016/j.tecto.2011.07.001>.
- Nemcok, M., Pospisil, L., Lexa, J., Donelick, R., 1998. Tertiary subduction and slab break-off model of the Carpathian-Pannonian region. *Tectonophysics* 295, 307–340. [https://doi.org/10.1016/S0040-1951\(98\)00092-4](https://doi.org/10.1016/S0040-1951(98)00092-4).
- Paffrath, M., Friederich, W., 2021. the AlpArray and AlpArray-Swath D working group. Imaging structure and geometry of slabs in the greater Alpine area – a P-wave traveltimes tomography using AlpArray seismic network data. *Solid Earth Discuss.*, 1–40. <https://doi.org/10.5194/se-2021-58>.
- Placer, L., 1998. Contribution to the macrotectonic subdivision of the border region between Southern Alps and External Dinarides. *Geologija* 41, 223–255. <https://doi.org/10.5474/geologija.1998.013>.
- Plomerová, J., Žlebčíková, H., Hetényi, G., Vecsey, L., Babuška, V., 2022. AlpArray-EASI working groups. Two subduction-related heterogeneities beneath the Eastern Alps and the Bohemian Massif imaged by high-resolution P-wave tomography. *Solid Earth* 13, 251–270. <https://doi.org/10.5194/se-13-251-2022>.
- Ratschbacher, L., Frisch, W., Linzer, H.G., Merle, O., 1991. Lateral extrusion in the Eastern Alps, part 2: structural analysis. *Tectonics* 10, 257–271. <https://doi.org/10.1029/90TC02623>.
- Roecker, S., Baker, B., McLaughlin, J., 2010. A finite-difference algorithm for full waveform teleseismic tomography. *Geophys. J. Int.* 181, 1017–1040. <https://doi.org/10.1111/j.1365-246X.2010.04553.x>.
- Royden, L.H., 1993. Evolution of retreating subduction boundaries formed during continental collision. *Tectonics* 12, 629–638. <https://doi.org/10.1029/92TC02641>.
- Schlunegger, F., Kissling, E., 2022. Slab load controls beneath the Alps on the source-to-sink sedimentary pathways in the Molasse Basin. *Geosciences* 12. <https://doi.org/10.3390/geosciences12060226>.
- Schmid, S.M., Bernoulli, D., Fügenschuh, B., Matenco, L., Schefer, S., Schuster, R., Tischler, M., Ustaszewski, K., 2008. The Alpine-Carpathian-Dinaridic orogenic system: correlation and evolution of tectonic units. *Swiss J. Geosci.* 101, 139–183. <https://doi.org/10.1007/s00015-008-1247-3>.
- Schmid, S.M., Pfiffner, O.A., Froitzheim, N., Schönborn, G., Kissling, E., 1996. Geophysical-geological transect and tectonic evolution of the Swiss-Italian Alps. *Tectonics* 15, 1036–1064. <https://doi.org/10.1029/96TC00433>.
- Schmid, S.M., Scharf, A., Handy, M.R., Rosenberg, C.L., 2013. The Tauern window (eastern Alps, Austria): a new tectonic map, with cross-sections and a tectonometamorphic synthesis. *Swiss J. Geosci.* 106, 1–32. <https://doi.org/10.1007/s00015-013-0123-y>.
- Schuster, R., Kurz, W., Krenn, K., Fritz, H., 2013. Introduction to the geology of the Eastern Alps. *Ber. Geol. Bundesanstalt Wien* 99, 121–133.
- Spada, M., Bianchi, I., Kissling, E., Agostinetti, N.P., Wiemer, S., 2013. Combining controlled-source seismology and receiver function information to derive 3-D Moho topography for Italy. *Geophys. J. Int.* 194, 1050–1068. <https://doi.org/10.1093/gji/ggt148>.
- Stüwe, K., Schuster, R., 2010. Initiation of subduction in the Alps: continent or ocean? *Geology* 38, 175–178. <https://doi.org/10.1130/G30528.1>.
- Sun, W., Zhao, L., Malusà, M.G., Guillot, S., Fu, L.Y., 2019. 3-D Pn tomography reveals continental subduction at the boundaries of the Adriatic microplate in the absence of a precursor oceanic slab. *Earth Planet. Sci. Lett.* 510, 131–141. <https://doi.org/10.1016/j.epsl.2019.01.012>.
- Tomek, C., Hall, J., 1993. Subducted continental margin imaged in the Carpathians of Czechoslovakia. *Geology* 21, 535–538. [https://doi.org/10.1130/0091-7613\(1993\)021<0535:SCMIIN>2.3.CO;2](https://doi.org/10.1130/0091-7613(1993)021<0535:SCMIIN>2.3.CO;2).
- Ustaszewski, K., Schmid, S.M., Fügenschuh, B., Tischler, M., Kissling, E., Spakman, W., 2008. A map-view restoration of the Alpine-Carpathian-Dinaridic system for the Early Miocene. *Swiss J. Geosci.* 101, 273–294. <https://doi.org/10.1007/s00015-008-1288-7>.
- Valasek, P., Mueller, S., Frei, W., Holliger, K., 1991. Results of NFP 20 seismic reflection profiling along the Alpine section of the European Geotraverse (EGT). *Geophys. J. Int.* 105, 85–102. <https://doi.org/10.1111/j.1365-246X.1991.tb03446.x>.
- Vrabec, M., Fodor, L., 2006. Late Cenozoic tectonics of Slovenia: structural styles at the northeastern corner of the Adriatic microplate. In: Pinter, N., Gyula, G., Weber, J., Stein, S., Medak, D. (Eds.), *The Adria Microplate: GPS Geodesy, Tectonics and Hazards*. Springer Netherlands, Dordrecht, pp. 151–168.
- Wortel, M., Spakman, W., 2000. Subduction and slab detachment in the Mediterranean-Carpathian region. *Science* 290, 1910–1917. <https://doi.org/10.1126/science.290.5498.1910>.
- Yan, Q.Z., Mechie, J., 1989. A fine structural section through the crust and lower lithosphere along the axial region of the Alps. *Geophys. J. Int.* 98, 465–488. <https://doi.org/10.1111/j.1365-246X.1989.tb02284.x>.
- Zahorec, P., Papčo, J., Pašteka, R., Bielik, M., Bonvalot, S., Braitenberg, C., Ebbing, J., Gabriel, G., Gosar, A., Grand, A., Götz, H.J., Hetényi, G., Holzrichter, N., Kissling, E., Marti, U., Meurers, B., Mrlina, J., Nogová, E., Pastorutti, A., Salaun, C., Scarponi, M., Sebera, J., Seoane, L., Skiba, P., Szűcs, E., Varga, M., 2021. The first pan-Alpine surface-gravity database, a modern compilation that crosses frontiers. *Earth Syst. Sci. Data* 13, 2165–2209. <https://doi.org/10.5194/essd-13-2165-2021>.
- Zhao, L., Paul, A., Guillot, S., Solarino, S., Malusà, M.G., Zheng, T., Aubert, C., Salimbeni, S., Dumont, T., Schwartz, S., et al., 2015. First seismic evidence for continental subduction beneath the Western Alps. *Geology* 43, 815–818. <https://doi.org/10.1130/G36833.1>.
- Zhao, L., Paul, A., Malusà, M.G., Xu, X., Zheng, T., Solarino, S., Guillot, S., Schwartz, S., Dumont, T., Salimbeni, S., et al., 2016. Continuity of the Alpine slab unraveled by high-resolution P wave tomography. *J. Geophys. Res.*, Solid 121, 8720–8737.