



OPEN ACCESS

EDITED BY

Nathalie Oulhen,
Brown University, United States

REVIEWED BY

Paul Yancey,
Whitman College, United States
Wanchun Guan,
Wenzhou Medical University, China

*CORRESPONDENCE

Silvia Simonetti

✉ silvia.simonetti2@unisi.it

RECEIVED 08 February 2025

ACCEPTED 18 April 2025

PUBLISHED 08 May 2025

CITATION

Simonetti S, Mutemi KN, Romano P, Luckenbach T, Zupo V, Gambi MC and Corsi I (2025) Hydrothermal vents as observatories for future ocean acidification (OA) scenarios: an *in-situ* study to unravel the involvement of ATP binding cassette transporters in the adaptation of marine polychaetes *Platynereis* spp. to OA. *Front. Mar. Sci.* 12:1573367. doi: 10.3389/fmars.2025.1573367

COPYRIGHT

© 2025 Simonetti, Mutemi, Romano, Luckenbach, Zupo, Gambi and Corsi. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Hydrothermal vents as observatories for future ocean acidification (OA) scenarios: an *in-situ* study to unravel the involvement of ATP binding cassette transporters in the adaptation of marine polychaetes *Platynereis* spp. to OA

Silvia Simonetti^{1,2*}, Kevin Nzumbi Mutemi³, Patrizia Romano^{1,4}, Till Luckenbach⁵, Valerio Zupo², Maria Cristina Gambi⁶ and Ilaria Corsi¹

¹Department of Physical, Earth and Environmental Sciences, University of Siena, Siena, Italy,

²Department of BluBioTech, Stazione Zoologica Anton Dohrn, National Institute of Marine Biology, Ecology and Biotechnology, Napoli, Italy, ³Developmental Biology Unit, European Molecular Biology Laboratory (EMBL), Heidelberg, Germany, ⁴Department of Biology and Evolution of Marine Organisms (BEOM), Stazione Zoologica Anton Dohrn, National Institute of Marine Biology, Ecology and Biotechnology, Napoli, Italy, ⁵Department Ecotoxicology, Helmholtz Centre for Environmental Research (UFZ), Leipzig, Germany, ⁶National Institute of Oceanography and Applied Geophysics (OGS), Trieste, Italy

The marine annelid *Platynereis dumerilii*, is a key model in genetics, evolution, neurobiology, ecology, and ecotoxicology. Along with its sibling species, *P. cfr massiliensis*, it thrives in both normal and naturally acidified environments. This makes these species ideal candidates for studying mechanisms of tolerance to acidified conditions, resembling future ocean acidification (OA) scenarios. The ATP-binding cassette (ABC) transport proteins help mitigating the adverse impacts of drugs, xenobiotics and physical stressors. There is growing evidence for their involvement to mediate tolerance towards acid-stress in bacteria and tumor cell lines. Such a function may be relevant for the ability of marine species to cope with OA and may be important to consider when predicting future OA scenarios for marine fauna. Here we addressed the question if ABC transporters of *Platynereis* spp. are involved in compensating adverse effects of low pH by studying ABC transporter transcript levels in marine animals exposed to various pH levels. We firstly examined *P. dumerilii* whole genome data (version EMBL_pдум_1.0, Genbank assembly: GCA_026936325.1) for the presence of ABC transporter genes, by homology searches, and, using the single-cell atlas database with *P. dumerilii* gene expression data, we then determined the presence of a potentially relevant subset of ABC transporters from the ABCB, C and G subfamilies in different organs/tissues. Finally, to assess how seawater pH affects ABC transporter expression, we conducted an *in-situ* reciprocal transplant experiment involving individuals of *P. dumerilii*/*P. cfr massiliensis*. Adult specimens were collected inside and outside the CO₂ vents

off Castello Aragonese (Ischia Island, Italy). Individuals collected from normal pH areas (8.18 ± 0.005) were transplanted to acidified conditions (7.33 ± 0.312), and *vice versa*, while others were placed in their original areas. We found 81 orthologs from ABC transporter subfamilies A-G, expressed in different organs/tissues including midgut, neurons, body epidermis and ectodermal cells, and somatic and visceral muscle. Following the 30 days transplant experiment, qPCR analyses were performed to examine the expression levels of seven selected genes from the ABCB, ABCC, and ABCG subfamilies (*abcb_1*, *abcb_2*, *abcb_3*, *abcc_1*, *abcc_2*, *abcc_3*, and *abcg*). Three of these genes were differentially expressed in specimens transplanted from normal pH to low pH areas (*abcb_1* and *abcg* up-regulated while *abcb_3* down-regulated). Based on the homology with human *ABCB1* and *ABCG2*, which are crucial in tumor cell adaptation to acidified environments, it seems reasonable to hypothesize that *abcb_1*, *abcb_3* and *abcg* play a similar role in *Platynereis* spp. helping in maintaining cellular homeostasis and surviving acid stress.

KEYWORDS

ocean acidification, CO₂ vents, *Platynereis* spp., ATP binding cassette (ABC) transporters, acid stress tolerance, gene expression analysis

1 Introduction

In certain marine ecosystems, CO₂ concentrations in the water are naturally elevated, which is due to gas emissions from submarine volcanism (vents) or from nutrient upwelling (González-Delgado and Hernández, 2018; Santana-Casiano et al., 2016). The pCO₂ in such waters can be several folds higher than the pCO₂ commonly found in oceanic surface waters (150–530 μatm) (Takahashi et al., 2014). Gradients of seawater pCO₂ levels in areas with natural CO₂ emissions, such as the volcanic CO₂ system in the Mediterranean Sea off Castello Aragonese on Ischia Island (Bay of Naples, Italy), can range from levels predicted for oceanic surface waters by the year 2100 due to climate change (500–1370 μatm, IPCC, 2014) to extreme conditions exceeding 20,800 μatm (Hall-Spencer et al., 2008). Studies of the species inhabiting such naturally acidified marine areas can provide insights into adaptations that enable organisms to tolerate lowered pH levels. This information can be useful to predict the effects of the globally growing anthropogenic ocean acidification (OA) on marine ecosystems. OA refers to the ongoing decrease in the surface seawater pH from a global, anthropogenic increase of the CO₂ concentration in seawater. This is related to the emissions of anthropogenically produced CO₂ into the atmosphere from combustion of fossil fuels that sharply increased after the industrial revolution and since then remained at a high level and are even still increasing (Bindoff et al., 2019; Takahashi et al., 2014).

Although negative effects of OA to marine life have been recognized, certain marine species that inhabit naturally acidified habitats are adapted to tolerate low pH conditions and thrive in them. Understanding the molecular/cellular bases of the adaptation

to hostile acidified conditions and the molecular/cellular bases of tolerance/resistance could unravel the consequences of future OA scenarios for the marine invertebrate fauna.

Various studies provided evidence that ATP-binding cassette (ABC) transporter proteins confer cellular tolerance in acid stress environments in bacteria and in tumor cells, although the mechanism how ABC transporters counteract the adverse effects of low pH to cells and tissue is not understood (Aye et al., 2009; Gao et al., 2023; Leslie et al., 2005; Matsushashi et al., 2015; Shvarev and Maldener, 2020; Tahara et al., 2012; 2015; Uchiyama et al., 2019; Wei and Roepe, 1994; Zhu et al., 2019, 2022). Some insights on these mechanisms were obtained from studies on cyanobacteria, as *Synechocystis* sp (Matsushashi et al., 2015; Tahara et al., 2012; 2015; Uchiyama et al., 2019) and *Anabaena* sp (Shvarev and Maldener, 2020), exposed to acidified environments (pH from 3.0 to 6.0). Other bacteria able to adapt to low pH environments, like fermented foods for *Acetobacter pasteurianus* (Gao et al., 2023) and *Lactococcus lactis* (Zhu et al., 2019; 2022), have been investigated. For all the above-mentioned species, survival in acidified environments has been attributed to the presence of ABC transporters that is likely to have a role in conferring resistance to adverse pH conditions. It was suggested that the tolerance to acid stress in bacteria is due to the ABC transport of substances crucial for the maintenance of homeostasis in cells, such as the export of lipids or negatively charged substances to regulate the extracellular structures as external barriers against H⁺ ions (Matsushashi et al., 2015; Tahara et al., 2012; 2015; Uchiyama et al., 2019) or the import of ions enhancing the activity of intracellular enzymes for the equilibration of internal pH (Zhu et al., 2022). Additional important findings on the involvement of ABC

transport proteins in acid stress tolerance/adaptation originated from studying Multidrug Resistance (MDR) and pH of cancer cell lines. Wei and Roepe (1994) have checked if lower extracellular pH, typical of malignant cells, contributes to the development of the MDR phenotype in colon carcinoma SW620 and HCT15 lines never exposed to chemotherapeutic drugs. Results have shown an increase in ABCB1/MDR1 protein content and gene expression in cells exposed to a CO₂-enriched external environment (pH reached the value of 6.5). The adaptation of the SW620 and HCT15 lines to acid external pH has been attributed to the increased expression and functional responses of MDR related ABC transporters that are probably associated with re-equilibration of the pH of the cytosol (as shown by Keizer and Joenje, 1989), facilitating the physiological functions of the cells. These findings indicate that ABC transporters can be involved in the adaptation to low pH conditions on the cellular level; however, it is so far not clear whether ABC transporters are involved in acid-stress resistance on the multicellular organism-level.

The ABC transporters are among the largest and well conserved protein superfamilies identified to date, comprising so far ten subfamilies (ABCA to J) across all prokaryote and eukaryote taxa (Dean and Annilo, 2005; Verrier et al., 2008). There has been great attention towards the ABCB, ABCC and ABCG subfamilies because of their involvement in MDR and Multixenobiotic Resistance (MXR), denominating cellular/organismal resistance towards environmental xenobiotics (Kurelec, 1992), and their importance in bacterial antibiotic resistance (Greene et al., 2018). Originally known for their involvement in cancer drug resistance (Biedler and Riehm, 1970), they are more generally a cellular stress response as the first and last defense system widely present in several taxa including marine species and their embryos/larvae stages (Epel et al., 2008; Kurelec, 1992). Beyond drugs and xenobiotics, MDR genes are induced by other stress signals, such as heat-shock, inflammation, hypoxia, UV and X irradiation (Scott, 2003).

Very limited information is currently available on the mechanisms of tolerance to low pH conditions of marine invertebrate species, despite their importance to understand the potential future impact of OA on marine life (Simonetti et al., 2022). The knowledge on bacterial and cancer cell resistance to acid stress provides a valuable foundation to address research on the topic. It is reasonable to hypothesize that ABC transporters may play a similar role in helping more complex species to tolerate low-pH conditions. Additionally, the mechanisms carried out by bacterial and cancer cells might also be employed by multicellular organisms to manage acidic conditions, aiding in the maintenance of homeostasis and the prevention of cell death.

The aim of the present study was to disclose the involvement of ABC proteins in adaptation to low-pH/high-pCO₂ environments in the marine polychaete sibling species *Platynereis dumerilii/cfr massiliensis* populating naturally acidified marine coastal vents. *Platynereis dumerilii* (Audouin and Milne-Edwards, 1834) is a marine annelid polychaete belonging to the family Nereididae, widely distributed throughout the European seas, all along the

Mediterranean coasts and the Black Sea either in normal and acidified environments (Read and Fauchald, 2023). It is considered a model species in genetics and single-cell genomics, because its genome has been completely sequenced, which provides an important base for molecular development and evolution studies (Özpolat et al., 2021; Tessmar-Raible and Arendt, 2003) in comparative studies since its evolutionary lineage has been slow-evolving (Zantke et al., 2014), and in neurobiology, ecology (Hardege, 1999) and ecotoxicology (Bellan, 1980; Hutchinson et al., 1995). Furthermore, it has been proven to be an excellent model to address various aspects of acclimation/adaptation to OA, such as functional traits analyses (Lucy et al., 2015; Gambi et al., 2016) and evolutionary implications (Wäge et al., 2017). It is highly abundant, along with its sibling species *P. cfr massiliensis*, in naturally acidified areas, such as in the CO₂ vents off Castello Aragonese (Ischia Island, Italy) (Calosi et al., 2013; Ricevuto et al., 2012; 2014; Wäge et al., 2018), as well as in other vent systems (e.g., Vulcano vents, Italy; Vizzini et al., 2017; Wäge et al., 2017).

Firstly, the presence of ABC genes as annotated sequences from the known genome and their expression in various organs/tissues of adult specimens of *P. dumerilii*. *cfr massiliensis* was investigated. Then, the contribution of 7 selected ABC transporter proteins to adaptation/tolerance to low-pH/high-pCO₂ seawater conditions in two annelid sibling species was established thanks to a 30 d *in-situ* reciprocal transplant experiment in a naturally acidified marine area characterized by vents with a seawater pH gradient from normal pH value (8.1) down to 6.5.

2 Materials and methods

2.1 Genome-wide identification of ABC transporters in *P. dumerilii*

2.1.1 Identification and characterization of putative ABCs

The presence of the ABC transport proteins in the genome of *P. dumerilii* was investigated in tissues and organs *via* homology searches as detailed below. The *P. dumerilii* genome version EMBL_pdum_1.0 (Genbank assembly: GCA_026936325.1) was used as the reference assembly for ABC transporter identification (Mutemi et al., 2024, biorxiv).

All the available protein sequences of animal ABC transporters were downloaded from the National Center for Biotechnology Information (NCBI) database (Supplementary Table S1) and a local alignment with the *P. dumerilii* proteome was conducted by the BLAST tool (Basic Local Alignment Search Tool). A Pfam analysis was also carried out by the hmmer tool, version 3.3.2 (Eddy, 2011), to catch any other potential ABC transport protein sequence not identified by the BLAST analysis. The most specific domains for ABC proteins (listed in Supplementary Table S2) from <https://pfam.xfam.org/> were selected and used for the analysis. Annotation and nomenclature of each *P. dumerilii* ABC protein were completed

based on amino acid sequence similarities in terms of *in silico* analysis by searching for known orthologs, amino acid similarity and identity comparison with BLASTp analysis on the protein database of NCBI. Finally, annotated proteins were searched in the *P. dumerilii* genome to disclose the corresponding coding genes.

2.1.2 Phylogenetic analysis of ABCs

The ABC protein amino acid sequences of *P. dumerilii* were phylogenetically analyzed to show the evolutionary relationships among them based upon similarities and differences in their characteristics. The full-length amino acid sequences retrieved from the NCBI server were aligned using T-Coffee web server (Notredame et al., 2000) and a phylogenetic tree was constructed with iTOL v6 web server (Letunic and Bork, 2021). Furthermore, the ABC protein sequences previously retrieved from NCBI (Supplementary Table S1) were used for the phylogenetic analysis with the *P. dumerilii* ABC proteins to highlight evolutionary relationships between species and to support the previous ABC annotations. For this, full-length amino acid sequences were aligned using the Muscle tool, version 5.1 (Edgar, 2022) and trimmed by the ClipKIT tool, version 1.3.0 (Steenwyk et al., 2020), using the most restrictive trimming method kpic-smart-gap to retain phylogenetically informative sites and remove the others. A phylogenetic tree was constructed with a maximum likelihood method using the iq-tree tool, version 2.2.0.3 (Nguyen et al., 2014).

2.1.3 Organ/tissue distribution and expression of ABC genes

In order to understand the possible physiological function of ABC transport proteins, their distribution in the *P. dumerilii* body was examined through the ‘P.Dumwhole-body’ single cell atlas available at EMBL Heidelberg. The cellular atlas integrates gene

expression and ultrastructure for the whole *P. dumerilii* body and cells are clustered in organ/tissue classes based on single-cell RNA sequencing.


A subset of 40 ABC transporter genes was selected from ABCB, C and G subfamilies and searched into the EMBL available atlas for their tissue distribution and expression pattern.

2.2 *In-situ* reciprocal transplant experiment

2.2.1 Study area

The transplant study was conducted at the volcanic vents off Castello Aragonese (40° 43′ 57.9″ N, 13° 57′ 51.8″ E), located on the north-eastern side of Ischia Island (Tyrrhenian Sea, Bay of Naples, Italy) (Figure 1). The naturally acidified area is characterized by thermal vents with a seawater pH gradient from normal value (8.1) down to 6.5 (0.7 x 10⁶ L d⁻¹ emitted gas at the north side and 1.4 x 10⁶ L d⁻¹ emitted gas at the south side, Hall-Spencer et al., 2008). In this area, three sites (A4, A5, A6; “A” for “Acidified”) were chosen to collect specimens of polychaetes and allocate the moorings for the *in-situ* reciprocal transplant experiment (Figure 1).

The three sites at normal pH (8.1) were chosen in the San Pietro area (40° 44′ 47.6″ N; 13°56′ 40.42″E), approximately 4 km off the vents. Indeed, the environmental characteristics of the area were very similar to those characterizing the vents in terms of exposure to light and hydrodynamic conditions, temperature, salinity, and depth, except for the pH value (Calosi et al., 2013). Specimens of polychaetes were collected nearby San Pietro, and three mooring stations (N1, N2, N3) were placed approximately 50 m from each other (Figure 1). A further site at normal pH was used for specimens’ collection due to limited abundance in the San Pietro area and named S. Anna (40° 43′ 34.51″ N, 13° 57′ 35.7″ E) located approximately 600 m away from the south side



of the vent area of Castello, no venting activities are present in this area and the pH is in the range between 8.05 and 8.11 (Ricevuto et al., 2015a) (Figure 1).

Specimens were also collected from the acidified area of Castello Aragonese and three mooring stations (A4, A5 on the south side, and A6 on the north side) were placed again approx. 50 m from each other (Figure 1).

Seawater pH, salinity and conductivity in each site were recorded, at the beginning and at the end of the transplant experiment, as follows: A4: pH 7.25 ± 0.03 , salinity 38.5 ± 0.71 ‰, conductivity 52.8 ± 0.07 $\mu\text{S mm}^{-1}$; A5: pH 7.62 ± 0.01 , salinity 39.0 ± 0.0 ‰, conductivity 53.0 ± 0.07 $\mu\text{S mm}^{-1}$; A6: pH 6.88, salinity 38.0 ± 0.0 ‰, conductivity 52.0 ± 0.14 $\mu\text{S mm}^{-1}$; N1-3: pH 8.18 ± 0.01 , salinity 38.5 ± 0.5 ‰, conductivity 53.0 ± 0.125 $\mu\text{S mm}^{-1}$. Long-term trend of the pH in these zones are also available in Foo et al. (2018).

2.2.2 Sample collection

Adult individuals of the two sibling species *P. dumerilii*/*P. cfr massiliensis* living in association with macroalgae, mainly the brown algae *Halopteris scoparia* and *Dictyota* spp., were collected during the first week of May 2023. Specimens of both species were collected by SCUBA divers at 1–2 m depth from the acidified area at the southern sites of Castello Aragonese and by snorkelers at 1–2 m depth from the two normal-pH sites, “San Pietro” and “S. Anna” rocks. Thalli of *H. scoparia* and *Dictyota* spp. with *Platynereis* individuals attached were removed and immediately transferred to containers filled with natural seawater on board and transported to the laboratory. The algae were then transferred to large plastic trays, to avoid anoxic conditions, and gently shaken to isolate *Platynereis* spp. specimens. After morphological identification, specimens were measured by means of millimetric paper and divided into three size groups (< 1 cm, 1–2 cm, > 2 cm length).

Organisms were maintained for 1–3 d in a temperature-controlled room at 18°C at a light regime of 12 h light/12 h dark in plastic bowls (10 individuals per bowl) supplied with approximately 200 mL of seawater taken from the sampling site to reduce the stress due to handling. pH was also checked daily and maintained constant at 8.2 for specimens collected from normal pH sites and between 7.2 and 7.6 for the acidified sites.

Additional 40 specimens from N and A areas were stored for molecular analysis of time 0 of the reciprocal transplant experiment (t0_N and t0_A groups respectively, Table 1) and then transferred into Eppendorf tubes with RNAlater solution (Sigma-Aldrich Company Ltd., Gillingham, U.K.).

2.2.3 Transplant of individuals

Individuals collected at sites with the normal seawater pH were placed in the moorings located in the acidified area and exposed for 30 d (May-June 2023) (t30_NA); other individuals were left in the moorings within the original site at normal seawater pH and left for 30 days as well (t30_N). Similarly, specimens collected from the acidified site were transferred in moorings placed in the normal-pH (t30_AN) area and some left in the acidified one and both left for 30 days (t30_A) in accordance with previous transplant studies (Ricevuto et al., 2015b) in which 30 days were selected to stimulate major pathway of response (Table 1). The experiment was designed to use two replicates of 10 individuals per each experimental plot (however, due to bad weather conditions, only one replicate was available for t30_NA4, t30_NA5, t30_NA6 treatments).

Moorings were made of cylindrical chambers hosting at least 10 individuals each and homogeneously divided according to their size. Two small cylindrical chambers were placed for each mooring. Transplantation chambers were constructed from PVC tubes (diameter 4 cm, length 11 cm) closed at both ends by a plankton net (mesh size 100 mm). The net mesh size guarantees regular water flow through the tube, allowing respiration and preventing escapes. These chambers have been used in previous experimental transplants (see Calosi et al., 2013). Small portions of *H. scoparia* and *Dictyota* spp. thalli were added to each chamber as a feeding source and a substrate to attach, to maintain polychaetes in conditions as close as possible to those they experience in their habitat. The chambers were fixed, via plastic cable ties, to a rope attached to stony moorings (Figure 2), and deployed by SCUBA divers in each exposure site, approximately at 2 m depth.

After 30 days, chambers were carefully retrieved by snorkeling and rapidly transferred to the laboratory where polychaetes were checked for healthy conditions and then stored at -80°C in RNAlater for subsequent analyses.

TABLE 1 Summary of sample IDs and experimental conditions.

Sample ID	Initial pH (mean)	Final pH (mean)	Transplant	Collection time
t0_A	7.25-7.62	–	–	t0 – before transplant
t0_N	8.18	–	–	t0 – before transplant
t30_A4	7.25-7.62	7.62	A → A4	t30 – end of transplant
t30_A5	7.25-7.62	7.25	A → A5	t30 – end of transplant
t30_A6	7.25-7.62	6.88	A → A6	t30 – end of transplant
t30_N	8.18	8.18	N → N	t30 – end of transplant
t30_AN	7.25-7.62	8.18	A → N	t30 – end of transplant
t30_NA4	8.18	7.62	N → A4	t30 – end of transplant
t30_NA5	8.18	7.25	N → A5	t30 – end of transplant
t30_NA6	8.18	6.88	N → A6	t30 – end of transplant




FIGURE 2

Stony moorings with anchored cylindrical pvc chambers where worms were allocated at the beginning (left) and 30 days later, at the end (right) of the experiment in one of the plots within the vents (A5).

A schematic description of the experimental plan is reported in [Figure 3](#).

2.3 Quantitative analysis of ABC transporter gene expression

2.3.1 Total RNA isolation and cDNA synthesis

Total RNA was extracted from pools of five *P. dumerilii* individuals (4–18 mg tissue in total) using the PureLink[®] RNA Mini Kit following the manufacturer's protocol (Ambion life technologies, USA). In brief, samples stored in RNAlater at -80°C were thawed, centrifuged (6,000 rcf, 2 min, 4°C) and the supernatant was removed. The tissue was homogenized using the TissueLyser LT (Qiagen) with a run time of 3 min at 50 oscillations. The integrity of the extracted total RNA was tested on a BioPhotometer (Eppendorf 6131) and on a 1.5% agarose 'bleach gel' ([Aranda et al., 2012](#), modified); the RNA concentration was quantified with a BioPhotometer (Eppendorf). First-strand cDNA

was synthesized from approximately 130 ng of RNA in a final volume of 20 µL by using the qScript Ultra SuperMix[®] (Quantabio, USA) according to manufacturer's instructions (8 µL RNA template, 4 µL qScript XLT cDNA SuperMix (5X), 8 µL RNase/DNase-free water).

2.3.2 Target gene selection

Based on previous studies on the involvement of ABCs in the resistance to acid conditions of bacteria species and cancer cells ([Gao et al., 2023](#); [Matsuhashi et al., 2015](#); [Shvarev and Maldener, 2020](#); [Tahara et al., 2012](#); 2015; [Uchiyama et al., 2019](#); [Wei and Roepe, 1994](#); [Zhu et al., 2019](#)), target genes belonging to the superfamily of ABC were selected from the *P. dumerilii* ABC inventory. Here the focus was on the subfamilies ABCB, ABCC, ABCG which are the largest subfamilies and in particular the *abcb* genes (*abcb_1*, *abcb_2*, *abcb_3*) were chosen for their homology to ABCB1/MDR1/P-gp and ABCB4 genes, *abcc* genes (*abcc_1*, *abcc_2*, *abcc_3*) were selected for the homology with MRPs and *abcg* gene is homologous to ABCG2 multidrug transporter ([Supplementary](#)




FIGURE 3

The worms' transplant experimental plan.

Table S3). Nucleotide sequences for the selected genes were obtained from *Platynereis* genome (version 2.0) sequenced at EMBL. The sequences of forward and reverse primers are reported in Supplementary Table S3 and were designed through the online tool Primer3Plus (Untergasser et al., 2007).

The two housekeeping genes, *18S* and *α-TUB* and respective primers (Supplementary Table S3) were from Wäge et al. (2018).

2.3.3 qPCR analysis of gene expression

qPCRs were performed in triplicate on an iCyclerIQ5 (Bio-Rad, USA). Reaction mixes with a final volume of 20 μL contained 2 μL cDNA (diluted 1:10), 10 μL PerfeCTa® SYBR® Green SuperMix, Low ROX (Quantabio, USA) and 1 μL (10 μM) of each primer pair. A reaction without cDNA template was run in parallel to control for primer dimers and/or external contamination. qPCR conditions were: initial denaturation at 95°C for 30 s, followed by 40 cycles of denaturation at 95°C for 5 s, annealing at 60°C for 15 s and extension at 72°C for 10 s. For the generation of a melt curve, 61 cycles of 10-s increasing temperature steps from 65°C to 95°C were added at the end of the qPCR run. The primer efficiency (Supplementary Table S3) was determined by a three-point cDNA dilution series (dilution factor of 5) and calculated by the iCyclerIQ5. The relative quantification of gene expressions was analyzed by the formula (Livak and Schmittgen, 2001).

$$(E * 0.01 + 1)^{-\Delta\Delta CT}$$

Where *E* corresponds to primer efficiency and *CT* represents the threshold cycle values.

2.4 Statistical analyses

For statistical analyses, data were tested for normal distribution using the Kolmogorov–Smirnov test. As normal distribution of data was not found for all cases, the significance of variations in transcription levels were evaluated using the nonparametric Kruskal–Wallis and Dunn's *post hoc* tests. For all analyses, a *p* < 0.05 was considered statistically significant. Values were presented as means ± standard deviations (SD) and significant differences were labeled with asterisks and letters. Statistical analyses of gene expression were performed using GraphPad Prism 5 for Windows (GraphPad Software, USA).

3 Results

3.1 Genome-wide identification of ABC transporters in *P. dumerilii*

3.1.1 Identification, characterization and phylogenetic analysis of *P. dumerilii* ABC transporters

A total of 81 ABC transporters were annotated and identified in *P. dumerilii* proteome and genome, belonging to the 7 subfamilies

from A to G, as shown in Figure 4 and listed in Supplementary Table S4.

The largest ABC, encoded by the gene XLOC_024852, was of 2583 amino acids and the shortest ones, encoded by XLOC_028182 and XLOC_033621 of 132 (Supplementary Table S4). Subfamily C was the largest, with 34 transport proteins, followed by subfamilies B and G with 14 proteins, while 12 transport proteins belonged to the subfamily A. Subfamilies D and F included 3 proteins each and the smallest subfamily resulted the ABCE, with only 1 protein (Supplementary Table S4). No proteins from the subfamily H were found.

Among the 81 putative ABC transport proteins identified, 33 proteins were full transporters (2 nucleotide-binding domains – NBDs and 2 transmembrane domains – TMDs), belonging to the subfamilies ABCA, ABCB and ABCC, while 25 were half transporters (1 NBD and 1 TMD), from the subfamilies ABCA, ABCB, ABCC, ABCD and ABCG (Supplementary Figure S1). One ABCC transporter contained 2 NBDs and 3 TMDs; three proteins being part of the subfamilies ABCA, ABCB and ABCC contained 2 NBDs and 1 TMD and three of the subfamilies ABCB and ABCG contained 1 NBD and 2 TMDs. In addition, 16 proteins with high similarity to ABC transport proteins from other organisms missed the NBD or the TMD. Those were considered as incomplete proteins, possibly due to the completion and annotation of the *P. dumerilii* proteome.

Phylogenetic analysis of *P. dumerilii* ABC transporter proteins and all the available protein sequences of ABC transporters from NCBI database confirmed the previous annotation (Figure 5).

3.1.2 Organ/tissue distribution of ABC genes expression in *P. dumerilii*


Thanks to the *P. Dum* whole-body single cell atlas, clusters of tissue-prevalent and tissue-specific genes and genes that are differentially expressed in different tissues/organs were identified as shown in Figure 6. Most of the ABC transport proteins found in *P. dumerilii* are midgut-specific (8) or midgut-prevalent (9). Some showed specificity for neurons (5), body epidermis and ectodermal cells (2), and somatic (1) and visceral (1) muscle. The remaining ABCs were prevalent in unknown-tissue clades (2) or broadly expressed in all tissues (12).

3.2 *In-situ* transplant experiment

At the end of the 30 days, mortality was observed only in t30_N groups (2.5%). All worms appeared in good health since they exhibited their protective mucous tubes and could actively move when taken out from chambers.

3.2.1 Differential ABC gene expression in organisms inhabiting different pH environments

As far as gene expression profile of ABC transport proteins in specimens from the transplant experiments, a comparison between specimens transplanted and those left in the original areas was



vary (Figure 9A). On the contrary, *abc1* gene expression resulted in down-regulation in specimens from A to N ($p < 0.01$) (Figure 9B).

abc2: similarly, this gene also resulted in up-regulation ($p < 0.05$) from t0_N group to t30_NA5 as well as t30_NA4, although not statistically significant. At lower pH (6.88, t30_NA6) no significant differences with t0_N were observed (Figure 9A). No changes in the *abc* gene expression were recorded in specimens translocated from A to N (Figure 9B).

abc3: a down-regulation was observed from t0_N to t30_NA5 ($p < 0.05$). Similarly, t30_NA6 specimens showed a decrease, but not significant, in *abc3* gene expression compared to t0_N. No significant differences were detected in specimens transferred from N to A4 compared to the t0_N treatment (Figure 9A). Interestingly, the reverse transplant A to N induced an up-regulation of the *abc3* gene ($p < 0.01$) (Figure 9B).

abc1: similarly, a down-regulation was observed from t0_N to t30_NA4 and t30_NA6 but only significant between t0_N and

t30_NA6 group ($p < 0.05$) (Figure 9A). Unfortunately, no data was available for the t30_NA5 treatment. The individuals collected at t0_A did not show any statistically significant differences in *abc1* gene expression with those transplanted from A to N (t30_AN) (Figure 9B).

abc2: no significant differences were found in t30_NA5 and t30_NA6 compared to t0_N while up-regulation of the *abc2* was recorded in t30_NA4 specimens compared to the t0_N ones ($p < 0.05$) (Figure 9A). *abc2* gene expression in treatment transplanted from A to N showed no statistically significant changes compared to t0_A (Figure 9B).

abc3: no significant changes in the expression of this gene were recorded (Figure 9).

abcg: an up-regulation was recorded in specimens from t0_N to t30_NA5 ($p < 0.05$) and between t0_N and t30_NA6 ($p < 0.01$) (Figure 9A). Contrarily, a down-regulation was observed in specimens translocated from A to N compared to t0_A group ($p < 0.01$) (Figure 9B).




FIGURE 6

Dotplot of ABC transport protein tissue/organ distribution and expression visualizing the relative expression levels of a subset of ABC transporter genes (x-axis) across all tissue/organ clusters (y-axis). Fold gene expression levels are indicated by shades of red (increase in gene expression) and blue (decrease in gene expression), and larger dot diameter indicates that the gene expression was detected in greater proportion of cells from the cluster.

4 Discussion

4.1 Genome-wide identification of ABC transporters in *P. dumerilii*

4.1.1 Identification, characterization and phylogenetic analysis of *P. dumerilii* ABC transporters

The *P. dumerilii* set of ABC transport proteins clearly is by far one of the largest with 81 proteins (Supplementary Table S5). From the available literature, only *Tetranychus urticae* and *Macrostomum lignano* have a higher number of ABC transporters (103 and 92 respectively) (Caña-Bozada et al., 2019; Dermauw et al., 2013), while from a minimum of 9 (*Neobenedenia melleni*, Caña-Bozada et al., 2019) to a maximum of 73 genes have been reported for other species (*Tribolium castaneum*, Broehan et al., 2013) (Supplementary Table S5). Furthermore, *P. dumerilii* shows a higher number of abc transporter proteins compared to marine Chordata species and *Homo sapiens* (Supplementary Table S5). Gene duplication has been identified as a mechanism for genomic adaptation to changing environments (Kondrashov, 2012). Therefore, the increased number of ABC transporter genes, as the one found in *P. dumerilii*, might be related to its ability to withstand

extreme conditions, including the CO₂ venting areas at Ischia Island (Calosi et al., 2013).

P. dumerilii ABC transport proteins can be divided into 7 subfamilies, labeled from A to G. This is consistent with our current understanding of ABC transporter subfamilies, as these seven subfamilies were found to be present across all taxa (Li and Nikaido, 2004; Annilo et al., 2006; Rea, 2007; Jeong et al., 2017). Notably, we observed the absence of subfamily H, which was identified in invertebrates and a single vertebrate species, the zebrafish (*Danio rerio*) (Dean and Annilo, 2005). Additionally, subfamily ABCI is found exclusively in plants (Verrier et al., 2008), while the subfamily ABCJ was recently proposed as a separate group, including nine proteins from the mosquito *Aedes aegypti* (Figueira-Mansur et al., 2020).

The ABCB, ABCC and ABCG subfamilies resulted in our study the most expanded, as observed among aquatic invertebrates (Jeong et al., 2017). Actually, these subfamilies include many highly conserved MXR-mediated related efflux transporters (abcb1/p-gp, abcc1-5, abcg2) which constitute important cellular defense mechanism towards marine pollutants in marine species (Epel, 1998; Kurelec, 1992). According to Xu et al. (2023), the number of ABCD, ABCE, and ABCF members was consistent across species, except for Platyhelminthes, which have a variable amount of such




FIGURE 7

Relative gene expression of the 7 selected ABC genes in *Platynereis* spp. specimens collected at time 0 in normal (N) (white bars) and naturally acidified (A) (black bars) sites. Results are shown as mean \pm standard deviation (SD). Asterisks indicate significant differences (* p < 0.05, ** p < 0.01) between the two groups.

genes. As such, this confirms a high level of conservation for these transport proteins.

Phylogenetic analysis revealed that ABCE and ABCF subfamilies were more closely related to each other than to the other subfamilies (Figure 5).

Based on phylogenetic analysis including the *P. dumerilii* ABC transporter proteins and all protein sequences of animal ABC transporters available on the NCBI database (Figure 5) each *P. dumerilii* ABC transporter has homologues within the ABC transporters of *H. sapiens* and *Mus musculus*, and in many cases of *Rattus norvegicus*. For a more detailed characterization of *P. dumerilii* ABC transport proteins subfamilies see [Supplementary Materials](#).

4.1.2 Organ/tissue distribution of ABC gene expression in *P. dumerilii*

ABC proteins from the subfamilies B, C and G have been discovered to be highly expressed in the midgut of different insects, including *Tribolium castaneum*, *Conogethes punctiferalis*, *Bactrocera dorsalis* and *Nilaparvata lugens*. Most of them have been shown to be involved in resistance/clearance of insecticides and insecticidal proteins (Li et al., 2020; Rösner and Merzendorfer, 2020; Shwe et al., 2022; Wang et al., 2023). These findings suggest that the abundance of ABC proteins belonging to the subfamilies b, c and g found in our study in the midgut and neurons of *P. dumerilii* may be related to their involvement in resistance towards toxicants as well. ABC proteins of the subfamilies b and g participate in the resistance of cancer cells to low-pH environments as well. Therefore, it is reasonable to assume that ABC proteins belonging to the subfamilies B and G might have an important role in adaptation to low pH in seawater also in *P. dumerilii* by playing a role in

physiological function maintenance and buffering the internal cell pH.

4.2 *In-situ* reciprocal transplant experiment

Stability of the ABC transporter gene expression was evaluated in specimens before the transplant and after 30 days of experiment in both normal- and low-pH areas as follows: t0_N vs t30_N and t0_A vs t30_A4, t30_A5, t30_A6 (Figure 8). Such assessment was needed to eventually exclude those genes that can be differentially expressed by other external factors beyond pH changes. All 7 ABC genes resulted stable except for *abcb_2* and *abcc_2* so these 2 were not selected for the next evaluation steps of their differential expression upon changes in pH. Hypotheses on the involvement of each *abc* gene in adaptation to low-pH were made based on the basal expression gene level of organisms inhabiting different pH environments (Figure 7) and verified through the ABC transcription results of the *in-situ* reciprocal transplant experiment (Figure 9).

abcb_1 was stably expressed across time in 2 out of 3 A sites, except in the t30_A5 vs t0_A (Figure 8B). Its expression was significantly different between the two populations inhabiting different pH habitats, with an increase in t0_A group compared to t0_N (Figure 7) so the enhancement of *abcb_1* gene transcription could contribute to the resistance of *Platynereis* spp. to low pH. The hypothesis was confirmed thanks to the results of the transplant experiment, in which it was up-regulated in specimens translocated to low pH sites (t30_NA5 and t30_NA4) (Figure 9A) as well as down-regulated in organisms reversely translocated from A to N (t30_AN) (Figure 9B). The *abcb_1* gene is homologous to the




FIGURE 8

Gene expression profile of ABC genes in specimens from (A) N sites at t0 (white bars) and after 30 days (t30) (grey bars) and (B) A sites at the beginning (t0) (white bars) and at the end (t30) (grey bars) of the experiment in the three plots (A4, A5, A6). Results are shown as mean \pm standard deviation (SD). Asterisks indicate significant differences ($*p < 0.05$, $**p < 0.01$, $***p < 0.001$) with the (A) t0_N group and (B) t0_A group.

human *ABCB1/MDR1* gene, found to be involved in tumor cell adaptation to acidified environments. Similarly to the *Platynereis* spp. *abcb_1*, the human *MDR1* gene was detected to be up-regulated in colon carcinoma SW620 and HCT15 cell lines exposed for 5 d to pH 6.5 (Wei and Roepe, 1994). Interestingly, when the cells were shifted back to a normal extracellular pH (7.0), the gene expression decreased, in agreement with that observed in

our study in specimens from A to N. Authors hypothesize that the increase in MDR protein transcription allows cells to cope with the acidic external environment by maintaining intracellular pH homeostasis (Wei and Roepe, 1994). Increased internal cell pH is correlated with the level of MDR expression by pumping acidic substances (Keizer and Joenje, 1989). Thus, it is reasonable to assume that *abcb_1* in *Platynereis* spp. plays an important role in




FIGURE 9

Relative gene expression levels of *ABC* genes in specimens transplanted from (A) N (white bars) to A4, A5, A6 (grey bars) and (B) A (white bars) to AN (grey bars). Results are shown as mean \pm standard deviation (SD). Asterisks indicate significant differences (* $p < 0.05$, ** $p < 0.01$) with the (A) t0_N group and (B) t0_A group.

adaptation to low pH in seawater by playing a similar role and buffering the internal cell pH. It is worth noting that at t30_NA6 (pH 6.88) the transcription of the *abcb_1* gene in transplanted *Platynereis* spp. specimens were at the same level as non-transplanted worms (Figure 9A). It could indicate an hormetic behavior in the *abcb_1* gene expression to decreasing pH

environments. The phenomenon of hormesis is defined as an adaptive dose-dependent response characterized by a biphasic pattern with a rise at lower pressure and an inhibition with increasing stress conditions (Calabrese and Baldwin, 2002). Thus, this suggests a clear initial involvement of the gene in resistance to low pH conditions based on the up-regulation of the *abcb_1* gene

and the ineffectiveness of this mechanism at severe pH conditions. So, different strategies are applied by the worms to survive in more severe acidified environments.

The *abcb_3* gene expression did not change over time (Figure 8) and its decrease is likely to be involved in *Platynereis* spp. adaptation to low-pH environments, since it was down-expressed in individuals living in sites A as compared to those inhabiting normal-pH areas (Figure 7). The thesis was validated following the *in-situ* transplant experiment. Specimens transferred from N to A (t30_NA5 and t30_NA6) showed a down-regulation of *abcb_3* gene (Figure 9A) while those transplanted from A to N sites displayed the up-regulation of the gene (Figure 9B). As *abcb_1*, also *abcb_3* showed homology with human *ABCB1/P-glycoprotein (P-gp)*. However, an opposite behavior of *abcb_1* and *abcb_3* gene expression was observed in this study supporting the hypothesis of a multifunctionality role of P-gp in aquatic invertebrates (Jeong et al., 2017). Actually, from the single cell analysis of *Platynereis* spp., the two *abcb* proteins exhibited specificity for different cell clades: *abcb_1* is mostly present in the epidermis, midgut epithelium and smooth muscle cells while *abcb_3* seems to occur more in neurons. Therefore, it is possible that they absolve different functions including the resistance to low pH environments.

The *abcc_1* gene was stable over time (Figure 8) and by comparing t0_N and t0_A groups a significant down-expression was detected (Figure 7), confirmed also in specimens transplanted from N to A (t30_NA4, t30_NA6) (Figure 9A). However, a potential involvement in low pH resistance is not supported because of the absence of modulation in specimens translocated from A to N (Figure 8B). Similarly, although stable, *abcc_3* gene did not vary in specimens from N to A (Figure 9A) and reverse (Figure 9B).

Interestingly, the *abcg* gene was highly expressed in specimens living in A compared to those from N (Figure 7) and up-regulated in those transplanted from N to A (Figure 9A). A down-regulation was observed in specimens transferred from A to N (Figure 9B) in agreement with the results from Cheng and To (2012), who observed an increase in the *ABCG2* (homologous to *Platynereis* spp. *abcg*) gene expression in low-pH resistant human colon carcinoma HCT-116 cells after an exposure to pH 5. *ABCG2* is involved in the efflux of physiological metabolites from cells (Leslie et al., 2005) and performs an anti-apoptotic function (Aye et al., 2009) so it is possible to suppose that the enhancement of *abcg* under acid stress might help cells to maintain their homeostasis and prevent death.

5 Conclusions

A first attempt to provide a complete inventory of the ABC transporters of the annelid *P. dumerilii* was carried out here; 81 ABC transporters belonging to the seven subfamilies ABCA-G were identified. These findings were crucial for the following investigation on the involvement of ABC transporter proteins in *Platynereis* spp. resistance/adaptation to acidified conditions.

In summary, 3 out of the 7 *abcb* transporters, namely *abcb_1*, *abcb_3* and *abcg*, turned out to have a role in adaptation to low-pH/high-*pCO*₂ environments of *Platynereis* spp. populating naturally acidified marine coastal areas. The homology with human *ABCB1* and *ABCG2*, which are involved in carcinoma cell resistance to external low pH, suggests that the three here identified *abcb* transporters play a similar role during an exposure to acidified environments participating in the maintenance of cellular homeostasis, buffering the internal pH of internal cell pH cells and preventing cell death in order to guarantee the proper functioning of cellular activities. This might be possible thanks to the trafficking of molecules and substances enrolled in these processes, such as the efflux of physiological metabolites from cells, by means of the three identified ABC transporter proteins. In future studies the assessment of function of these proteins requires more attention in order to understand why the transporters represent an adaptation mechanism to acidified conditions.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

Ethics statement

The manuscript presents research on animals that do not require ethical approval for their study.

Author contributions

SS: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Validation, Visualization, Writing – original draft, Writing – review & editing. KM: Data curation, Formal Analysis, Investigation, Methodology, Software, Validation, Writing – review & editing. PR: Investigation, Writing – review & editing. TL: Validation, Writing – review & editing. VZ: Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Validation, Writing – review & editing. MG: Conceptualization, Funding acquisition, Investigation, Methodology, Project administration, Resources, Supervision, Validation, Writing – review & editing. IC: Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Validation, Writing – review & editing.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. This study was supported

by the 2020 scholarship co-founded by the University of Siena and the Stazione Zoologica Anton Dohrn of Naples entitled 'Molecular bases and evolutionary constraints of cellular acid tolerance in cyanobacteria and invertebrates subjected to ocean acidification and other sources of stress' (METASTRESS) in the framework of the Doctoral Research Program in Environmental, Geological and Polar Sciences and Technologies (36th cycle) of the University of Siena. Furthermore, part of the work was performed in the framework of the Erasmus Long Life Learning Program of Dr. Simonetti S. at EMBL Heidelberg.

Acknowledgments

We are grateful to Prof. Detlev Arendt for hosting SS during her Erasmus traineeship at EMBL Heidelberg. We also acknowledge Victoria Alicia Witte and Dr. Leslie Pan from the Developmental Biology Unit at EMBL Heidelberg for their contributions to the bioinformatics analysis. Additionally, we thank Prof. Maria Cristina Fossi's Lab (Department of Physical, Earth, and Environmental Sciences, University of Siena, Italy), particularly Dr. Cristina Panti and Dr. Giacomo Limonta, for their support with the real-time PCR instrument. We are also grateful to the Marine Protected Area staff "Regno di Nettuno" of Ischia for support at sea during the transplant experiment. At the beginning of this study, part of the PhD project of SS, MG was senior researcher at the Stazione Zoologica Anton Dohrn of Naples (Ischia Marine Center).

References

- Annilo, T., Chen, Z.-Q., Shulenin, S., Costantino, J., Thomas, L., Lou, H., et al. (2006). Evolution of the vertebrate ABC gene family: analysis of gene birth and death. *Genom.* 88, 1–11. doi: 10.1016/j.ygeno.2006.03.001
- Aranda, P. S., LaJoie, D. M., and Jorczyk, C. L. (2012). Bleach gel: A simple agarose gel for analyzing RNA quality. *Electrophoresis* 33, 366–369. doi: 10.1002/elps.201100335
- Aye, I. L., Singh, A. T., and Keelan, J. A. (2009). Transport of lipids by ABC proteins: interactions and implications for cellular toxicity, viability and function. *Chem. Biol. Interact.* 180, 327–339. doi: 10.1016/j.cbi.2009.04.012
- Bellan, G. (1980). Relationship of pollution to rocky substratum polychaetes on the French Mediterranean coast. *Mar. Pollut. Bull.* 1, 318–321. doi: 10.1016/0025-326X(80)90048-X
- Biedler, J. L., and Riehm, H. (1970). Cellular resistance to actinomycin D in Chinese hamster cells *in vitro*: cross-resistance, radioautographic, and cytogenetic studies. *Cancer. Res.* 30, 1174–1184.
- Bindoff, N. L., Cheung, W. W. L., Kairo, J. G., Aristegui, J., Guinder, V. A., Hallberg, R., et al. (2019). "Changing ocean, marine ecosystems, and dependent communities," in *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. Eds. H.-O. Pörtner, D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria, M. Nicolai, A. Okem, J. Petzold, B. Rama and N. M. Weyer (Cambridge University Press, Cambridge, UK and New York, NY, USA), 447–587. doi: 10.1017/9781009157964.007
- Broehan, G., Kroeger, T., Lorenzen, M., and Merzendorfer, H. (2013). Functional analysis of the ATP-binding cassette (ABC) transporter gene family of *Tribolium castaneum*. *BMC Genom.* 14, 6. doi: 10.1186/1471-2164-14-6
- Calabrese, E. J., and Baldwin, L. A. (2002). Defining hormesis. *Hum. Exp. Toxicol.* 21, 91–97. doi: 10.1191/0960327102ht217oa
- Calosi, P., Rastrick, S. P. S., Lombardi, C., de Guzman, H. J., Davidson, L., Jahnke, M., et al. (2013). Adaptation and acclimatization to ocean acidification in marine ectotherms: an *in situ* transplant experiment with polychaetes at a shallow CO₂ vent system. *Phil. Trans. R. Soc. Lond. B. Biol. Sci.* 368, 20120444. doi: 10.1098/rstb.2012.0444
- Caña-Bozada, T. V., Morales-Serna, F. N., García-Gasca, A., Llera-Herrerac, R., and Fajer-Ávila, E. J. (2019). Genome-wide identification of ABC transporters in monogeneans. *Mol. Biochem. Parasitol.* 234, 111234. doi: 10.1016/j.molbiopara.2019.111234
- Cheng, G. M., and To, K. K. (2012). Adverse cell culture conditions mimicking the tumor microenvironment upregulate ABCG2 to mediate multidrug resistance and a more Malignant phenotype. *ISRN Oncol.* 2012, 746025. doi: 10.5402/2012/746025
- Dean, M., and Annilo, T. (2005). Evolution of the ATP-binding cassette (ABC) transporter superfamily in vertebrates. *Annu. Rev. Genomics Hum. Genet.* 6, 123–142. doi: 10.1146/annurev.genom.6.080604.162122
- Dermauw, W., Osborne, E. J., Clark, R. M., Grbić, M., Tirry, L., and Van Leeuwen, T. (2013). A burst of ABC genes in the genome of the polyphagous spider mite *Tetranychus urticae*. *BMC Genom.* 14, 317. doi: 10.1186/1471-2164-14-317
- Eddy, S. R. (2011). Accelerated profile HMM searches. *PLoS Comput. Biol.* 7, e1002195. doi: 10.1371/journal.pcbi.1002195
- Edgar, R. C. (2022). Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence homology and phylogeny. *Nat. Commun.* 13, 6968. doi: 10.1038/s41467-022-34630-w
- Epel, D. (1998). Use of multidrug transporters as first lines of defense against toxins in aquatic organisms. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 120, 23–28. doi: 10.1016/S1095-6433(98)10005-3
- Epel, D., Luckenbach, T., Stevenson, C. N., MacManus-Spencer, L. A., Hamdoun, A., and Smital, T. (2008). Efflux transporters: newly appreciated roles in protection against pollutants. *Environ. Sci. Technol.* 42, 3914–3920. doi: 10.1021/es087187v
- Figueira-Mansur, J., Schrago, C. G., Salles, T. S., Alvarenga, E. S. L., Vasconcellos, B. M., Melo, A. C. A., et al. (2020). Phylogenetic analysis of the ATP-binding cassette proteins suggests a new ABC protein subfamily J in *Aedes aegypti* (Diptera: Culicidae). *BMC Genom.* 21, 463. doi: 10.1186/s12864-020-06873-8
- Foo, S. A., Byrne, M., Ricevuto, E., and Gambi, M. C. (2018). The carbon dioxide vents of Ischia, Italy, a natural laboratory to assess impacts of ocean acidification on

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The author(s) declared that they were an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmars.2025.1573367/full#supplementary-material>

- marine ecosystems: an overview of research and comparisons with other vent systems. *Ocean. Mar. Biol. Ann. Rev.* 56, 237–310.
- Gambi, M. C., Musco, L., Giangrande, A., Badalamenti, F., Micheli, F., and Kroeker, K. J. (2016). Distribution and functional traits of polychaetes in a CO₂ vent system: winners and losers among closely related species. *Mar. Ecol. Prog. Ser.* 550, 121–134. doi: 10.3354/meps11727
- Gao, L., Shi, W., and Xia, X. (2023). Genomic plasticity of acid-tolerant phenotypic evolution in *Acetobacter pasteurianus*. *Appl. Biochem. Biotechnol.* 195, 6003–6019. doi: 10.1007/s12010-023-04353-9
- González-Delgado, S., and Hernández, C. (2018). The importance of natural acidified systems in the study of ocean acidification: what have we learned? *Adv. Mar. Biol.* 80, 57–99. doi: 10.1016/bs.amb.2018.08.001
- Greene, N. P., Kaplan, E., Crow, A., and Koronakis, V. (2018). Antibiotic resistance mediated by the MacB ABC transporter family: A structural and functional perspective. *Front. Microbiol.* 9. doi: 10.3389/fmicb.2018.00950
- Hall-Spencer, J., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., et al. (2008). Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* 454, 96–99. doi: 10.1038/nature07051
- Hardege, J. D. (1999). Nereidid polychaetes as model organisms for marine chemical ecology. *Hydrobiologia* 402, 145–161. doi: 10.1023/A:1003740509104
- Hutchinson, T. H., Jha, A. N., and Dixon, D. R. (1995). The polychaete *Platynereis dumerilii* (Audouin and Milne-Edwards): a new species for assessing the hazardous potential of chemicals in the marine environment. *Ecotoxicol. Environ. Saf.* 31 (3), 271–81. doi: 10.1006/eesa.1995.1074
- IPCC. (2014). In Pachauri, R. K., and Meyer, L. A. (Eds.), *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva, Switzerland, 151.
- Jeong, C.-B., Kim, H.-S., Kang, H.-M., and Lee, J.-S. (2017). ATP-binding cassette (ABC) proteins in aquatic invertebrates: Evolutionary significance and application in marine ecotoxicology. *Aquat. Toxicol.* 185, 29–39. doi: 10.1016/j.aquatox.2017.01.013
- Keizer, H. G., and Joenje, H. (1989). Increased cytosolic pH in multidrug-resistant human lung tumor cells: effect of verapamil. *J. Natl. Cancer. Inst.* 81, 706–709. doi: 10.1093/jnci/81.9.706
- Kondrashov, F. A. (2012). Gene duplication as a mechanism of genomic adaptation to a changing environment. *Proc. R. Soc B* 279, 5048–5057. doi: 10.1098/rspb.2012.1108
- Kurelec, B. (1992). The multidrug resistance mechanism in aquatic organisms. *Crit. Rev. Toxicol.* 22, 23–43. doi: 10.3109/10408449209145320
- Leslie, E. M., Deeley, R. G., and Cole, S. P. C. (2005). Multidrug resistance proteins: role of P-glycoprotein, MRP1, MRP2, and BCRP (ABCG2) in tissue defense. *Toxicol. Appl. Pharmacol.* 204, 216–237. doi: 10.1016/j.taap.2004.10.012
- Leticun, I., and Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nuc. Acid. Res.* 49, W293–W296. doi: 10.1093/nar/gkab301
- Li, Z., Cai, T., Qin, Y., Zhang, Y., Jin, R., Mao, K., et al. (2020). Transcriptional response of ATP-binding cassette (ABC) transporters to insecticide in the brown planthopper, *Nilaparvata lugens* (Stål). *Insects* 11, 280. doi: 10.3390/insects11050280
- Li, X. Z., and Nikaido, H. (2004). Efflux-mediated drug resistance in bacteria. *Drugs* 64, 159–204. doi: 10.2165/00003495-200464020-00004
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* 25, 402–408. doi: 10.1006/meth.2001.1262
- Lucey, N. M., Lombardi, C., De Marchi, L., Schulze, A., Gambi, M. C., and Calosi, P. (2015). To brood or not to brood: are marine invertebrates that protect their offspring more resilient to ocean acidification? *Sci. Rep.* 5, 12009. doi: 10.1038/srep12009
- Matsuhashi, A., Tahara, H., Ito, Y., Uchiyama, J., Ogawa, S., and Ohta, H. (2015). Slr2019, lipid A transporter homolog, is essential for acidic tolerance in *Synechocystis* sp. PCC6803. *Photosynth. Res.* 125, 267–277. doi: 10.1007/s11120-015-0129-6
- Mutemi, K. N., Simakov, O., Pan, L., Santangeli, L., Null, R., Handberg-Thorsager, M., et al. (2024). A genome resource for the marine annelid *Platynereis dumerilii*. *BioRxiv*, 2024.06.21.600153. doi: 10.1101/2024.06.21.600153
- Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., and Minh, B. Q. (2014). IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* 32, 268–274. doi: 10.1093/molbev/msu300
- Notredame, C., Higgins, D. G., and Heringa, J. (2000). T-Coffee: A novel method for fast and accurate multiple sequence alignment. *J. Mol. Biol.* 302, 205–217. doi: 10.1006/jmbi.2000.4042
- Özpolat, B. D., Rande, N., Williams, E. A., Bezars-Calderón, L. A., Andreatta, G., Balavoine, G., et al. (2021). The Nereid on the rise: *Platynereis* as a model system. *EvoDevo* 12, 10. doi: 10.1186/s13227-021-00180-3
- Rea, P. A. (2007). Plant ATP-binding cassette transporters. *Annu. Rev. Plant Biol.* 58, 347–375. doi: 10.1146/annurev.arplant.57.032905.105406
- Read, G., and Fauchald, K. (2023). “World polychaeta database,” in *Platynereis dumerilii* (Audouin & Milne Edwards 1833).
- Ricevuto, E., Benedetti, M., Regoli, F., Spicer, J. I., and Gambi, M. C. (2015b). Antioxidant capacity of polychaetes occurring at a natural CO₂ vent system: Results of an *in situ* reciprocal transplant experiment. *Mar. Environ. Res.* 112, 44–51. doi: 10.1016/j.marenvres.2015.09.005
- Ricevuto, E., Kroeker, F., Ferrigno, F., Micheli, F., and Gambi, M. C. (2014). Spatio-temporal variability of polychaete colonization at volcanic CO₂ vents indicates high tolerance to ocean acidification. *Mar. Biol.* 161, 2909–2919. doi: 10.1007/s00227-014-2555-y
- Ricevuto, E., Lorenti, M., Patti, F. P., Scipione, M. B., and Gambi, M. C. (2012). Temporal trends of benthic invertebrate settlement along a gradient of ocean acidification at natural CO₂ vents (Tyrrhenian Sea). *Biol. Mar. Mediterr.* 19, 49–52.
- Ricevuto, E., Vizzini, S., and Gambi, M. C. (2015a). Ocean acidification effects on stable isotope signatures and trophic interactions of polychaete consumers and organic matter sources at a CO₂ shallow vent system. *J. Exp. Mar. Biol. Ecol.* 468, 105–117. doi: 10.1016/j.jembe.2015.03.016
- Rösner, J., and Merzendorfer, H. (2020). Transcriptional plasticity of different ABC transporter genes from *Tribolium castaneum* contributes to diflubenzuron resistance. *Insect Biochem. Mol. Biol.* 116, 103282. doi: 10.1016/j.ibmb.2019.103282
- Santana-Casiano, J. M., Fraile-Nuez, E., González-Dávila, M., Baker, E. T., Resing, J. A., and Walker, S. L. (2016). Significant discharge of CO₂ from hydrothermalism associated with the submarine volcano of El Hierro Island. *Sci. Rep.* 6, 25686. doi: 10.1038/srep25686
- Scotto, K. W. (2003). Transcriptional regulation of ABC drug transporters. *Oncogene* 22, 7496–7511. doi: 10.1038/sj.onc.1206950
- Shvarev, D., and Maldener, I. (2020). The HlyD-like membrane fusion protein All5304 is essential for acid stress survival of the filamentous cyanobacterium *Anabaena* sp. PCC 7120. *FEMS Microbiol. Lett.* 367, fnaa108. doi: 10.1093/femsle/fnaa108
- Shwe, S. M., Prabu, S., Jing, D., He, K., and Wang, Z. (2022). Synergistic interaction of Cry1Ah and Vip3Aa19 proteins combination with midgut ATP-binding cassette subfamily C receptors of *Conogethes punctiferalis* (Guenée) (Lepidoptera: Crambidae). *Int. J. Biol. Macromol.* 213, 871–879. doi: 10.1016/j.ijbiomac.2022.06.019
- Simonetti, S., Zupo, V., Gambi, M. C., Luckenbach, T., and Corsi, I. (2022). Unraveling cellular and molecular mechanisms of acid stress tolerance and resistance in marine species: New frontiers in the study of adaptation to ocean acidification. *Mar. Pollut. Bull.* 185, 114365. doi: 10.1016/j.marpolbul.2022.114365
- Steenwyk, J. L., Buida, T. J. III, Li, Y., Shen, X.-X., and Rokas, A. (2020). ClipKIT: A multiple sequence alignment trimming software for accurate phylogenomic inference. *PLoS Biol.* 18, e3001007. doi: 10.1371/journal.pbio.3001007
- Tahara, H., Matsuhashi, A., Uchiyama, J., Ogawa, S., and Ohta, H. (2015). Slr0751 and Slr1041 are involved in acid stress tolerance in *Synechocystis* sp. PCC 6803. *Photosynth. Res.* 125, 233–242. doi: 10.1007/s11120-015-0153-6
- Tahara, H., Uchiyama, J., Yoshihara, T., Matsumoto, K., and Ohta, H. (2012). Role of Slr1045 in environmental stress tolerance and lipid transport in the cyanobacterium *Synechocystis* sp. PCC6803. *Biochim. Biophys. Acta* 1817, 1360–1366. doi: 10.1016/j.bbabi.2012.02.035
- Takahashi, T., Sutherland, S. C., Chipman, D. W., Goddard, J. G., Ho, C., Newberger, T., et al. (2014). Climatological distributions of pH, pCO₂, total CO₂, alkalinity, and CaCO₃ saturation in the global surface ocean, and temporal changes at selected locations. *Mar. Chem.* 164, 95–125. doi: 10.1016/j.marchem.2014.06.004
- Tessmar-Raible, K., and Arendt, D. (2003). Emerging systems: between vertebrates and arthropods, the Lophotrochozoa. *Curr. Opin. Genet. Dev.* 13, 331–340. doi: 10.1016/S0959-437X(03)00086-8
- Uchiyama, J., Itagaki, A., Ishikawa, H., Tanaka, Y., Kohga, H., Nakahara, A., et al. (2019). Characterization of ABC transporter genes, *slr1180*, *slr1181*, and *slr1270*, involved in acid stress tolerance of *Synechocystis* sp. PCC 6803. *Photosynth. Res.* 139, 325–335. doi: 10.1007/s11120-018-0548-2
- Untergasser, A., Nijveen, H., Rao, X., Bisseling, T., Geurts, R., and Leunissen, J. A. M. (2007). Primer3Plus, an enhanced web interface to Primer3. *Nucleic Acids Res.* 35, W71–W74. doi: 10.1093/nar/gkm306
- Verrier, P. J., Bird, D., Burla, B., Dassa, E., Forestier, C., Geisler, M., et al. (2008). Plant ABC proteins – a unified nomenclature and updated inventory. *Trends Plant Sci.* 13, 151–159. doi: 10.1016/j.tplants.2008.02.001
- Vizzini, S., Martínez-Crego, B., Andolina, C., Massa-Gallucci, A., Connell, S. D., and Gambi, M. C. (2017). Ocean acidification as a driver of community simplification via the collapse of higher-order and rise of lower-order consumers. *Sci. Rep.* 7, 4018. doi: 10.0.4.14/s41598-017-03802-w
- Wäge, J., Rotchell, J. M., Gambi, M.-C., and Hardege, J. D. (2018). Target gene expression studies on *Platynereis dumerilii* and *Platynereis* cfr *massiliensis* at the shallow CO₂ vents off Ischia, Italy. *Estuar. Coast. Shelf Sci.* 207, 351–358. doi: 10.1016/j.ecss.2017.11.012
- Wäge, J., Valvassori, G., Hardege, J. D., Schulze and, A., and Gambi, M. C. (2017). The sibling polychaetes *Platynereis dumerilii* and *Platynereis massiliensis* in the Mediterranean Sea: are phylogeographic patterns related to exposure to ocean acidification? *Mar. Biol.* 164, 199. doi: 10.1007/s00227-017-3222-x
- Wang, L., He, W., Lu, J.-M., Sun, J., Jiang, S.-D., Wang, J.-J., et al. (2023). Characterization and transcriptional expression of ABCG genes in *Bactrocera dorsalis*: Insights into their roles in fecundity and insecticidal stress response. *Int. J. Biol. Macromol.* 253, 126836. doi: 10.1016/j.ijbiomac.2023.126836
- Wei, L.-Y., and Roepe, P. D. (1994). Low external pH and osmotic shock increase the expression of human MDR protein. *Biochem.* 33, 7229–7238. doi: 10.1021/bi00189a027

Xu, J., Zheng, J., Zhang, R., Wang, H., Du, J., Li, J., et al. (2023). Identification and functional analysis of ABC transporter genes related to deltamethrin resistance in *Culex pipiens pallens*. *Pest. Manage. Sci.* 79, 3642–3655. doi: 10.1002/ps.7539

Zantke, J., Bannister, S., Rajan, V. B. V., Raible, F., and Tessmar-Raible, K. (2014). Genetic and genomic tools for the marine annelid *Platynereis dumerilii*. *Genetics* 197, 19–31. doi: 10.1534/genetics.112.148254

Zhu, Z., Yang, Y., Yang, P., Wu, Z., Zhang, J., and Du, G. (2019). Enhanced acid-stress tolerance in *Lactococcus lactis* NZ9000 by overexpression of ABC transporters. *Microb. Cell. Factories*. 18, 136. doi: 10.1186/s12934-019-1188-8

Zhu, Z., Yang, P., Yang, J., and Zhang, J. (2022). Comparative transcriptome analysis reveals the contribution of membrane transporters to acid tolerance in *Lactococcus lactis*. *J. Biotechnol.* 357, 9–17. doi: 10.1016/j.jbiotec.2022.08.006