

**Laura AIROLDI, BALLESTEROS E., BUONUOMO R., VAN BELZEN J., BOUMA T.J., CEBRIAN E., DE CLERK O., ENGELEN A.H., FERRARIO F., FRASCHETTI S., GIANNI F., GUIDETTI P., IVESA L., MANCUSO F.P., MICHELI F., PERKOL-FINKEL S., SERRAO E.A., STRAIN E.M., MANGIALAJO L.**  
Dipartimento di Scienze Biologiche, Geologiche ed Ambientali, University of Bologna,  
Via S. Alberto 163, 48123, Ravenna Italy.  
Email: [laura.airoldi@unibo.it](mailto:laura.airoldi@unibo.it)

## **MARINE FORESTS AT RISK: SOLUTIONS TO HALT THE LOSS AND PROMOTE THE RECOVERY OF MEDITERRANEAN CANOPY-FORMING SEAWEEDS**

### **Abstract**

*Along Mediterranean coasts, canopy-forming seaweeds used to form diverse, productive and valuable “forest” habitats, but in the past decades conspicuous declines, sometimes to local extinction, have been reported in many regions. Canopies are retracting particularly close to urban areas, and are replaced by turf-forming and ephemeral algae or barrens. The persisting forests are under continued threat, and current protection measures are insufficient. We provide evidence that declines of canopy algae are dramatically extensive, and are driven by multiple local (nutrient enrichment and high sediment loads, fishing, heavy metal pollution) and global stressors (increasing temperature, high wave exposure). We also show that the combined management of local stressors (such as nutrients and sediments) would increase significantly the resilience of canopy algae to future climatic stressors, preventing their further deterioration. Finally, we discuss restoration prospects in areas where these systems have been lost. We conclude identifying the main needs to understand, guide and motivate effective conservation actions in these valuable ecosystems.*

**Key words:** Canopy algae, habitat loss, Mediterranean Sea, multiple threats, conservation

### **Introduction**

Along Mediterranean coasts, canopy-forming seaweeds (most frequently brown algae belonging to the order Fucales) form diverse, productive and valuable “forest” habitats. These habitats are becoming rare at local, regional and basin scales at an alarming rate (Airoldi & Beck, 2007). This is concerning because algal canopies play a key role in coastal primary production and nutrient cycling, and facilitate rich flora and fauna communities. In the past decades, algal canopies have suffered widespread and apparently irreversible loss, much of which may have gone unnoticed. Algal canopies are retracting particularly close to urban areas, and are replaced by turf-forming and ephemeral algae or sea urchin barrens, with major negative consequences for associated benthic and fish communities (Benedetti-Cecchi *et al.*, 2001). The persisting forests are under continued threat, and the benefits of current protection measures have been low.

We synthesize past research efforts aiming at quantifying the losses, and identifying what factors drive the loss or enhance the resilience of these systems. We also discuss the restoration prospects in areas where canopies have been lost and the main needs.

### **Materials and methods**

We reviewed published primary literature and summarized it in a table. The review is organized into three sections: 1) a compilation of data on historical loss of canopies along

Mediterranean coastlines and main drivers of loss; 2) a compilation of data on known factors enhancing resilience and restoration success; 3) a discussion of gaps in the data, ecological knowledge, and protection measures for these coastal habitats and recommendations for how to address these gaps.


## Results

### Historical loss and main drivers

Conspicuous declines of algal canopies, sometimes to local extinction, have been reported in many regions along the coasts of Spain, France, Italy, Croatia, Albania, Greece and Turkey (Fig. 1 and Tab. 1). Along the Albères coast only 5 out of 14 species of Fucales (*Cystoseira* spp. and *Sargassum* spp.) documented as abundant in 1912 were present in 2003 (Thibaut *et al.*, 2005). Lost algal forests tend to be replaced by assemblages of lower structural complexity, such as turf-forming, filamentous or other ephemeral seaweeds, mussels or “barrens” (Mangialajo *et al.*, 2008; Connell *et al.*, 2014; Strain *et al.*, 2014). Canopy algae, turfs and barrens have been suggested to represent alternative states in shallow temperate rocky coasts under different disturbance and stress regimes (Airoidi *et al.*, 2009). There is a growing consensus and empirical evidence that these habitat shifts are driven by multiple anthropogenic stressors, including overfishing of higher trophic groups leading to outbreaks of grazers, eutrophication, excess sediment loads, coastal development, heavy metal pollution, point source pollutants such as oil spills, detergents and anti-fouling paints and invasive species (Table 1). These local anthropogenic stressors can interact negatively with environmental stressors or global climatic stressors (such as increasing temperature and CO<sub>2</sub>) resulting in accelerated declines of canopy-algae (Perkol-Finkel & Airoidi, 2010; Asnaghi *et al.*, 2013; Olabarria *et al.*, 2013; Strain *et al.*, 2014).

### Factors enhancing resilience or restoration efforts

While the proximate drivers of canopy loss are now relatively well understood, the factors that control the recovery have been more difficult to identify, and over a certain deterioration threshold, these systems may not be able to recover at all (Perkol-Finkel & Airoidi, 2010). The alternative habitat replacing lost canopies seems



**Fig. 1: Canopy algae characterise Mediterranean rocky coasts (e.g. A-*Cystoseira balearica* forest, Scandola, Corse), but many forests have been replaced by algal turfs (e.g. B- Haifa, Israel), urchin barrens (e.g. C- Porto Cesareo, Italy), or mussel beds (e.g. D - Monte Conero, Italy). Photographs by: A) E Ballesteros, B) L Airoidi, C) P Guidetti, D) L Airoidi**

to play a crucial role in controlling the return to a canopy dominated state once the stressors have been removed (Airoldi *et al.*, 2009). Thus, while there are examples of recovery of canopies from outbreak of urchins when effective protection measures have been established (*e.g.* Guidetti, 2006), when algal forests become replaced by turfs, sediments, or mussels it is not yet clear what prevents the recovery of the system, other than severe recruitment failure (Perkol-Finkel & Airoldi, 2010). Recent work has shown that management of water quality and particularly sediment loads is critical for ensuring the persistence of *Cystoseira* forests (Sales *et al.* 2011, Strain *et al.* unpub.). Reduction of nutrients would provide the greatest opportunity to prevent the shift from canopy to turf algae because of the prevalence of synergistic interactions between nutrients with other local and global stressors (Strain *et al.*, 2014). If depletion of forests has already occurred over wide areas, natural recovery could be slow or even implausible (Perkol-Finkel & Airoldi, 2010). The artificial restoration of fucoids in the Mediterranean Sea has been understudied compared to kelps and fucoids in other parts of the world, but results so far suggest it could be an effective strategy (Sales *et al.*, 2011; Perkol-Finkel *et al.*, 2012; Gianni *et al.*, 2013).

### **Discussion**

Mediterranean canopy forests are affected by many threats. The greatest impacts are associated with degraded water quality, coastal development, outbreaks of herbivores and invasive species, while effects of diseases and climate changes are uncertain. Current losses are alarming and protection is insufficient. Some key needs and opportunities for conservation and management are suggested below:

1) There is no comprehensive summary of the distribution of canopy forests, particularly deep sea ones, and their management is impeded by lack of knowledge on their status. Detailed habitat mapping should be given priority. The ecosystem services that these coastal habitats provide (such as nurseries for fisheries and recreation) also need to be better assessed to illustrate the costs of their loss and provide impetus and economic incentives for their protection and restoration.

2) An adequate evolutionary framework is needed to inform decisions on local and regional species diversity and to differentiate local extinction from species extinction. The loss of genetic biodiversity as populations undergo bottlenecks is also undescribed. New molecular tools need to be applied (RAD-seq) or developed (SNPs or microsatellites) to assess genetic diversity and link it to population resilience and ecosystem functioning, assess connectivity of populations, and study parentage and recruitment at local scales.

3) Like in other ecosystems (*e.g.* lakes, coral reefs, or forests) a gradual degradation of resilience paves the loss of these algal forests to alternative habitats, so that the mere restoration of environmental conditions preceding the loss may be insufficient to restore the system (Perkol-Finkel & Airoldi, 2010). Strategies for conservation of canopy forests should focus on “early-warning signals” of approaching shifts and on effective and rapid management of local stressors to maintain resilience in face of global stressors. This knowledge is presently limited for Mediterranean canopy forests, but results so far suggest that management of water quality and sediment loads would provide some of the greatest opportunities, particularly in enclosed bays or estuaries.

**Tab. 1: Selected studies reporting the loss (as either percentage area lost and/or number of species lost) of canopy-forming algae in the Mediterranean Sea, suggested drivers of loss, and signs of recovery if observed. Na= no quantitative data reported**

Lost species	Geographic location	Amount of loss	Driver of loss	Alternative habitat	Signs of recovery	References
<i>Cystoseira</i> spp. and <i>Sargassum acinarium</i> (as <i>S. linifolium</i> )	Monte Conero (Italy, Central Adriatic Sea)	90% (6 out of 8 sp lost)	High sediment loads/poor water quality, increased substratum instability, increased storminess	Turfs or mussels	no	(Romagnoli & Solazzi, 2003; Irving <i>et al.</i> , 2009; Perkol-Finkel & Airoldi, 2010)
<i>Cystoseira</i> spp. and <i>Sargassum</i> spp	Albères Coast (France, NW Mediterranean)	80% (9 out of 14 sp lost)	Poor water quality, overgrazing by urchins, coastal development, human trampling	Turfs, mussels or urchin barrens	no	(Thibaut <i>et al.</i> , 2005)
<i>Cystoseira</i> spp. and <i>Fucus virsoides</i>	Istrian coast (Croatia, North Adriatic Sea)	11 out of 15 sp lost	Poor water quality, overgrazing by urchins	Ephemeral algae	yes	(Munda, 1979, 2000; Zavodnik <i>et al.</i> , 2002)
<i>Cystoseira</i> spp	Genova (Italy, Ligurian Sea)	>50%	Coastal urbanisation	Algal turfs	no	(Mangialajo <i>et al.</i> , 2008)
<i>Cystoseira</i> spp	Albania	90%	Coastal development and urbanisation			(Fraschetti <i>et al.</i> , 2011)
<i>Cystoseira</i> spp. and <i>Sargassum</i> spp	Tremiti islands (Italy, South Adriatic Sea)	na	Poor water quality, overgrazing by urchins	Ephemeral algae or urchin barrens	yes	(Cormaci & Furnari, 1999; Fraschetti <i>et al.</i> , 2012)
<i>Cystoseira</i> spp	Salento Peninsula, (South Adriatic and Ionian Sea)	90%	Overgrazing by urchins	Urchin barrens	yes	(Guidetti <i>et al.</i> , 2003; Guidetti, 2006)
<i>Cystoseira</i> spp and <i>Sargassum</i> spp	Linosa Island (Italy, Sicily Channel)	60% (10 out of 15 sp lost)	Increased water temperature and changes in water circulation	Turfs and ephemeral algae	no	(Serio <i>et al.</i> , 2006)
<i>Cystoseira</i> spp	Several locations (Italy, NW Mediterranean)	90%	Coastal urbanisation, poor water quality, high sediment loads	Turfs	no	(Benedetti-Cecchi <i>et al.</i> , 2001)
<i>Cystoseira</i> spp	Several locations, (Greece and Turkey, Southern Aegean Sea)	>90%	Invasive fish	Urchin-less barrens	no	(Sala <i>et al.</i> , 2011; Giakoumi, 2014)
<i>Cystoseira crinita</i>	Cap Corse (France, NW Mediterranean)	7%	Coastal development	Turfs	no	(Sales & Ballesteros, 2010)
<i>Cystoseira crinita</i> , <i>C. barbata</i>	Maó harbour, Menorca, (Spain, Balearic Sea)	100%	Poor water quality, heavy metal pollution, and aquaculture	Ephemeral algae	Yes after transplant	(Sales <i>et al.</i> , 2011)
<i>Cystoseira zosteroides</i>	Medes Islands, (Spain, NW Mediterranean)	90%	Exceptional storm	Turfs	yes	(Navarro <i>et al.</i> , 2011)

4) Fucoids show high reproductive potential but low dispersal, which limits their natural recovery of wide lost/degraded areas. Given the extent of damage, restoration will be required in many places to meet any reasonable goals for conservation and management. Artificial restoration of *Cystoseira* forests in the Mediterranean Sea is much behind compared to other systems (i.e. seagrass beds), and much more work is needed to develop effective tools and approaches (Gianni *et al.*, 2013).

There are still opportunities for conservation of Mediterranean canopy forests. This protection should be achieved quickly because conservation is cheaper than restoration. Reducing cumulative local human impacts would represent the most effective strategy for the conservation and recovery of these systems, but, whenever this alone cannot reverse the loss, well-designed restoration projects can assist. Overall, there should be greater public, political and even scientific awareness of the extent, importance, and consequences of the loss of canopy forests, and greater commitment to motivate serious conservation and restoration actions in these highly threatened ecosystems.

### Acknowledgments

The study was supported by project TETRIS (Prin 2010-2011, MIUR).

### Bibliography

- AIROLDI L., BECK M.W. (2007) - Loss, status and trends for coastal marine habitats of Europe. *Oceanogr. Mar. Biol. Annu. Rev.*, 45: 345-405.
- AIROLDI L., CONNELL S.D., BECK M.W. (2009) - The loss of natural habitats and the addition of artificial substrata. In: M. Wahl (ed), *Marine hard bottom communities: patterns, dynamics, diversity, and change*, Springer, New York: 269-280.
- ASNAGHI V., CHIANTORE M., MANGIALAJO L., GAZEAU F., FRANCOUR P., ALLIOUANE S., GATTUSO J.P. (2013) - Cascading effects of ocean acidification in a rocky subtidal community. *PLoS ONE*, 8.
- BENEDETTI-CECCHI L., PANNACCIULLI F., BULLERI F., MOSCHELLA P.S., AIROLDI L., RELINI G., CINELLI F. (2001) - Predicting the consequences of anthropogenic disturbance: large-scale effects of loss of canopy algae on rocky shores. *Mar. Ecol. Prog. Ser.*, 214: 137-150.
- CONNELL S.D., FOSTER M.S., AIROLDI L. (2014) - What are algal turfs? Towards a better description of turfs. *Mar. Ecol. Prog. Ser.*, 495: 299-307.
- CORMACI M., FURNARI G. (1999) - Changes of the benthic algal flora of the Tremiti Islands (southern Adriatic) Italy. *Hydrobiologia*, 399: 75-79.
- FRASCHETTI S., BEVILACQUA S., GUARNIERI G., TERLIZZI A. (2012) - Idiosyncratic effects of protection in a remote marine reserve. *Mar. Ecol. Prog. Ser.*, 466: 21-34.
- FRASCHETTI S., TERLIZZI A., GUARNIERI G., PIZZOLANTE F., D'AMBROSIO P., MAIORANO P., BEQIRAJ S., BOERO F. (2011) - Effects of unplanned development on marine biodiversity: A lesson from Albania (central Mediterranean Sea). *J. Coast. Res.*: 106-115.
- GIAKOUMI S. (2014) - Distribution patterns of the invasive herbivore *Siganus luridus* (Ruppell, 1829) and its relation to native benthic communities in the central Aegean Sea, Northeastern Mediterranean. *Mar. Ecol.-Evol. Persp.*, 35: 96-105.
- GIANNI F., BARTOLINI F., AIROLDI L., BALLESTEROS E., FRANCOUR P., GUIDETTI P., MEINESZ A., THIBAUT T., MANGIALAJO L. (2013) - Conservation and restoration of marine forests in the Mediterranean Sea and the potential role of Marine Protected Areas. *Advances in Oceanography and Limnology*, 4: 83-101.
- GUIDETTI P. (2006) - Marine reserves reestablish lost predatory interactions and cause community changes in rocky reefs. *Ecol. Appl.*, 16: 963-976.



- GUIDETTI P., FRASCHETTI S., TERLIZZI A., BOERO F. (2003) - Distribution patterns of sea urchins and barrens in shallow Mediterranean rocky reefs impacted by the illegal fishery of the rock-boring mollusc *Lithophaga lithophaga*. *Mar. Biol.*, 143: 1135-1142.
- IRVING A.D., BALATA D., COLOSIO F., FERRANDO G.A., AIROLDI L. (2009) - Light, sediment, temperature, and the early life-history of the habitat-forming alga *Cystoseira barbata*. *Mar. Biol.*, 156: 1223-1231.
- MANGIALAJO L., CHIANTORE M., CATTANEO-VIETTI R. (2008) - Loss of furoid algae along a gradient of urbanisation, and structure of benthic communities. *Mar. Ecol. Prog. Ser.*, 358: 63-74.
- MUNDA I.M. (1979) - Some Fucacean association from the vicinity of Rovinj, Northern Adriatic. *Nova Hedwigia*, 31: 607-666.
- MUNDA I.M. (2000) - Long-term marine floristic changes around Rovinj (Istrian coast, North Adriatic) estimated on the basis of historical data from Paul Kuckuck's field diaries from the end of the 19th century. *Nova Hedwigia*, 71: 1-36.
- NAVARRO L., BALLESTEROS E., LINARES C., HEREU B. (2011) - Spatial and temporal variability of deep-water algal assemblages in the Northwestern Mediterranean: The effects of an exceptional storm. *Estuar. Coast. Shelf Sci.*, 95: 52-58.
- OLABARRIA C., ARENAS F., VIEJO R.M., GESTOSO I., VAZ-PINTO F., INCERA M., RUBAL M., CACABELOS E., VEIGA P., SOBRINO C. (2013) - Response of macroalgal assemblages from rockpools to climate change: effects of persistent increase in temperature and CO<sub>2</sub>. *Oikos*, 122: 1065-1079.
- PERKOL-FINKEL S., AIROLDI L. (2010) - Loss and recovery potential of marine habitats: an experimental study of factors maintaining resilience in subtidal algal forests at the Adriatic sea. *PLoS ONE*, 5: e10791.
- PERKOL-FINKEL S., FERRARIO F., NICOTERA V., AIROLDI L. (2012) - Conservation challenges in urban seascapes: promoting the growth of threatened species on marine coastal infrastructures. *J. Appl. Ecol.*, 49: 1457-1466.
- ROMAGNOLI T., SOLAZZI A. (2003) - Evoluzione dei popolamenti fitobentonici lungo la riviera del Conero dal 1941 al 2000. *Quad. Ist. Ric. Pesca Marittima, n.s.*, 1:63-84
- SALA E., KIZILKAYA Z., YILDIRIM D., BALLESTEROS E. (2011) - Alien marine fishes deplete algal biomass in the Eastern Mediterranean. *Plos One*, 6.
- SALES M., BALLESTEROS E. (2010) - Long-term comparison of algal assemblages dominated by *Cystoseira crinita* (Fucales, Heterokontophyta) from Cap Corse (Corsica, North Western Mediterranean). *Eur. J. Phycol.*, 45: 404-412.
- SALES M., CEBRIAN E., TOMAS F., BALLESTEROS E. (2011) - Pollution impacts and recovery potential in three species of the genus *Cystoseira* (Fucales, Heterokontophyta). *Estuar. Coast. Shelf Sci.*, 92: 347-357.
- SERIO D., ALONGI G., CATRA M., CORMACI M., FURNARI G. (2006) - Changes in the benthic algal flora of Linosa Island (Straits of Sicily, Mediterranean Sea). *Bot. Mar.*, 49: 135-144.
- STRAIN E.M., THOMSON R.J., MICHELI F., MANCUSO F.P., AIROLDI L. (2014) - Interactions of multiple stressors in driving the global loss of forests of canopy-forming algae to turf-forming algae and opportunities for remediation at local scale. *Glob. Change Biol.*, in press (DOI: 10.1111/gcb.12619)
- THIBAUT T., PINEDO S., TORRAS X., BALLESTEROS E. (2005) - Long-term decline of the populations of Fucales (*Cystoseira* spp. and *Sargassum* spp.) in the Alberes coast (France, North-western Mediterranean). *Mar. Poll. Bull.*, 50: 1472-1489.
- ZAVODNIK N., IVEŠA L., TRAVIZI A. (2002) - Note on recolonisation by furoid algae *Cystoseira* spp. and *Fucus virsoides* in the North Adriatic Sea *Acta. Adriat.*, 43: 25-32.