



Offshore fresh (or freshened) groundwater: Research achievements and future perspectives

Claudia Bertoni^{1,2,3} · Ariel Thomas⁴ · Hiba Wazaz⁴ · Angelo Camerlenghi³ · Jean-Christophe Comte⁵ · Davide Gamboa^{6,7} · Michela Giustiniani³ · Renée Martin-Nagle^{8,9} · Aaron Micallef¹⁰ · Margarita Nieves¹¹ · Elizabeth Quiroga Jordan¹² · Isha Savani^{3,12}

Received: 15 January 2026 / Accepted: 23 February 2026
© The Author(s) 2026

Abstract

Offshore fresh or freshened groundwater (OFG) is an important but under-represented component of coastal hydrogeological and climate systems. Recent advances in geophysical imaging, offshore drilling, numerical modelling, and machine learning have improved understanding of OFG distribution, dynamics, and connectivity with onshore aquifers. Remaining challenges include volumetric quantification, recharge history, ecological implications, and governance. OFG may complement coastal water-resource portfolios, but sustainable utilisation requires integrated management, environmental safeguards, and updated legal frameworks at national and international levels.

Keywords Groundwater and society · Groundwater exploration · Transboundary aquifers · Offshore fresh and freshened groundwater · Geophysical methods

Introduction

Offshore fresh (or freshened) groundwater, i.e. OFG (Post et al. 2013; Micallef et al. 2021), is an important and overlooked component of hydrological and climate frameworks, and its quantification can help refine water-availability

forecasts and support adaptation strategies in vulnerable coastal regions. The extent of OFG (Fig. 1), often reaching several kilometres beyond the coastline, indicates long-term storage of freshwater and variable connectivity with coastal aquifers (Arévalo-Martínez et al. 2023). These connections influence groundwater resilience to sea-level rise and salinisation, yet remain poorly constrained in current models. The question of whether the OFG component of the global water cycle should be considered in future climate-change models

Published in the special issue “Advances in understanding offshore freshened groundwater”.

✉ Claudia Bertoni
claudia.bertoni@earth.ox.ac.uk

¹ Department of Mathematics, Informatics and Geosciences, University of Trieste, Via Weiss, 2, 34128 Trieste, Italy

² Department of Earth Sciences, University of Oxford, South Parks Road, Oxford OX1 3AN, UK

³ National Institute of Oceanography and Applied Geophysics (OGS), 34010, Sgonico, Trieste, Italy

⁴ Department of Geosciences, University of Malta, 37 Triq Ta' Xmiexi, Msida, Malta

⁵ School of Geosciences, University of Aberdeen, St. Mary's Building, Elphinstone Road, Aberdeen AB24 3UF, UK

⁶ CESAM, University of Aveiro, Campus Universitário de Santiago, 3810-052 Aveiro, Portugal

⁷ Department of Geosciences, University of Aveiro, Campus Universitário de Santiago, 3810-052 Aveiro, Portugal

⁸ A Ripple Effect, 600 West Highland Avenue, Ebensburg, PA 15931, USA

⁹ Environmental Law Institute, 1730 M St NW #700, Washington, DC 20036, USA

¹⁰ Monterey Bay Aquarium Research Institute, 7700 Sandholdt Road, Moss Landing, CA 95039, USA

¹¹ University of Groningen, PO Box 72, 9700 AB Groningen, The Netherlands

¹² Ruden AS., Oslo Science Park, Gaustadalleen 21, 0349 Oslo, Norway




Fig. 1 Updated global map of OFG occurrences. Databases from: Post et al. (2013), Micallef et al. (2021), Giustiniani et al. (2025). Note the location of OFG studies that are part of this special issue

(SI): Chiacchieri et al. (2025); Corradin et al. (2025); Horozal et al. (2025); Li et al. (2025); Lin et al. (2025); Rahman et al. (2025); Sola et al. (2025)

(e.g. Waliser et al. 2007; Roberts et al. 2018), together with the implications for projections of coastal hydrology, and the quantification of the associated feedback. Groundwater resilience to sea-level rise, droughts, and salinisation yet remain poorly constrained in current models.

OFG can also have a significant and dynamic correlation with seafloor geomorphology, primarily through sediment destabilisation and associated fluid discharge processes (Hoffmann et al. 2023; Micallef et al. 2023; Saadatkhah et al. 2023). It influences marine ecology by facilitating biogeochemical fluxes from land to sea through submarine groundwater discharge (SGD, Taniguchi et al. 2002), a process that provides substantial fluxes of nutrients and carbon to coastal waters at levels comparable to major rivers (Moore 2010; Santos et al. 2008).

Understanding these environmental interactions is critical not only from an Earth system science perspective, but also for assessing OFG's viability and sustainability as a freshwater resource for drought-affected and water-stressed coastal regions (Bakken et al. 2012; UN-Water 2020). A fundamental consideration in this assessment is whether OFG systems contain fossil groundwater, emplaced during past sea-level lowstands and effectively non-renewable, or are actively recharged by present-day SGD. Consequently, in parallel with scientific efforts to identify and quantify OFG resources, attention has increasingly turned to their economic feasibility (Bakken et al. 2012) and associated legal frameworks (Martin-Nagle 2020), as well as the potential impacts of OFG abstraction on onshore groundwater

resources (Yu and Michael 2019). Beyond assessing utilization feasibility and direct impacts, these early evaluations offer a unique opportunity to establish foundational principles for the prospective development, management, and protection of this currently untapped resource, while also accounting for potential wider social and environmental impacts.

Nevertheless, comprehensive regional assessments of OFG remain uncommon, as they require either time-consuming, resource-intensive sub-seafloor data acquisition techniques or advanced integrated onshore-offshore 3D modelling. Lack of data leads to challenges in parameter selection in the models, which rarely include the required spatial complexity and the representation of long-term paleoenvironmental and paleoclimatic conditions. Recently, improvements in remote sensing technologies, subsea and subseafloor data acquisition tools and numerical methods have provided means to advance OFG investigations and shed light on the occurrence and extent of OFG. Emerging acquisition techniques now allow more efficient and cost-effective subsurface data collection. Simultaneously, the integration of multidisciplinary datasets and the adoption of advanced analytical approaches are substantially expanding our capacity to identify and characterize complex OFG systems. In this essay, we evaluate the current understanding of OFG in view of recent research, by highlighting unresolved questions and methodological challenges, and outlining emerging data acquisition and modelling methods, and research opportunities.

Timeline of OFG research

Although occurrences of deep, low-salinity or fresh groundwater within offshore sedimentary sequences have been noted earlier in unpublished oil & gas (O&G) well reports (Ruden 2007), the phenomenon only acquired scientific awareness following a drilling campaign along the U.S. continental margin (Kohout 1964; Hathaway et al. 1979). Terminology has varied considerably since its discovery (Table S1 of the electronic supplementary material (ESM)).

As interest in the subject expanded, the need for a unified and unambiguous term became apparent, leading to the adoption of the term OFG as “offshore fresh groundwater” (Post et al. 2013; Knight et al. 2018) or “offshore freshened groundwater” (Micallef et al. 2021). However, referring to these systems as “offshore” introduces an implicit spatial separation from onshore groundwater systems that does not necessarily exist from a hydrogeological perspective. In many settings, so-called offshore aquifers are directly connected to their onshore counterparts, forming a single groundwater system. Additionally, the use of the descriptor ‘freshened’ is often debated, as while it simply refers to salinity lower than ambient seawater (Micallef et al. 2021), for other authors the term suggests that groundwater was originally saline and it had been later diluted, in contrast with the widely accepted emplacement mechanisms of OFG as meteoric freshwater recharged during sea level lowstands.

Recently, in Europe, the COST Action CA21112 (OFF-SOURCE 2024) supported the creation of the first scientific network on OFG. The European Partnership Water4All funded the RESCUE project (RESCUE 2026) to build knowledge on deep-coastal and offshore low-salinity aquifers in the Northern Adriatic and North Sea. On a global scale, the International Ocean Drilling Programme (IODP³), and the National Science Foundation (NSF), with the European Consortium for Ocean Research Drilling (ECORD) as operator, implemented the first scientific drilling campaign dedicated entirely to OFG, Expedition 501 New England Shelf Hydrogeology (IODP³, 2025), which successfully validated a large OFG system hypothesis off the New England Shelf. Finally, the JPI Oceans launched a Joint Action ‘Offshore Freshened Groundwater’ (JPI Oceans 2025) to explore this untapped potential across European marine basins.

Established and emerging scientific methods

Since the Micallef et al. (2021) comprehensive review on OFG, several methodological advances have substantially expanded OFG investigation capabilities. On the geophysical side, new marine controlled-source electromagnetic (CSEM) hardware and survey concepts, such as the low-cost,

surface-towed system and novel onshore–offshore CSEM geometries, have broadened spatial coverage and reduced acquisition costs (Ishizu and Ogawa 2021; Pastorella et al. 2023). Additionally, trans-dimensional Bayesian inversion of CSEM data now yields probabilistic resistivity and salinity estimates instead of single deterministic models (Faghif et al. 2024).

Increasingly, 3-D hydrogeological models can integrate high-resolution seismic and sequence-stratigraphic analyses, allowing for geologically representative variable-density flow simulations (Thomas et al. 2022; De Biase et al. 2023; Campo et al. 2024). In parallel, multi-borehole porewater and isotope datasets, especially in deltaic and shelf settings, are being tightly coupled to transient paleo-hydrogeological models (Sheng et al. 2023), while 3D and global numerical modelling efforts now explore OFG dynamics, sustainability, and geomorphic impacts under changing sea-level and climate boundary conditions (Yu et al. 2025; Gupta and Micallef 2025; Zamrsky et al. 2022).

Limited offshore drilling, sparse geophysical imaging, and incomplete integration of onshore–offshore hydrogeological data still hinder robust assessments. However, borehole archives and methods from O&G exploration are currently being repurposed to compensate for this deficit and support regional assessments of OFG occurrence (Lipparini et al. 2023). Oil and gas (O&G) borehole data can provide direct evidence of low-salinity groundwater, provided that uncertainty is appropriately quantified and logs are subjected to rigorous petrophysical reinterpretation for hydrogeological purposes (Quiroga et al. 2023). On the other hand, O&G geophysical data such as seismic reflection data can be mostly used to define the presence of suitable reservoirs for OFG (Bertoni et al. 2020).

The use of O&G data remains limited when such datasets have commercial value and are covered by confidentiality agreements, which is commonly the case. Notable exceptions are national open-access repositories maintained by governmental agencies, which publish information derived from regional O&G activities, such as UK, Italy, Norway, Australia and New Zealand’s national data repositories (e.g. NDR n.d.); VIDEPI repository n.d.).

Emerging methods that could also help address the current uncertainties include machine learning (ML), which offers multiple ways to enhance the characterisation of OFG by leveraging advances established in subsurface and groundwater sciences (Pourghasemi et al. 2020). A key application is in the analysis of legacy O&G data, as automated raster digitisation of legacy well logs, geological sections, and maps greatly accelerate the creation of large, standardised databases, reducing the labour-intensive manual work traditionally required (Katole et al. 2025). Additionally, methods including neural networks and unsupervised clustering have demonstrated strong performance

in geological mapping (Cracknell and Reading 2014), offering transferable workflows for delineating offshore aquifers. In hydrogeology, supervised and unsupervised ML models have been widely applied to groundwater quality assessment, aquifer properties, vulnerability, recharge, and groundwater-level prediction (Adombi et al. 2021). The applications to OFG research have been so far much more limited (Haffert et al. 2024; Thomas et al. 2025). However, it is recognised that implementing ML techniques to determine OFG occurrences has the potential to overcome some of the constraints imposed by traditional methodological approaches. ML frameworks guided by domain expertise are better suited for use in OFG research to complement, rather than replace, process-based models by diagnosing structural mismatches and reducing uncertainty in predictions.

Impacts, economy and environment: Utilisation and challenges

Assessing OFG's resource potential requires evaluating its competitiveness relative to established alternatives such as desalination and conventional freshwater supplies. This requires analysing several key parameters, including the quality of the water, extraction and transport costs, environmental impacts, and treatment requirements. In an early assessment of extraction feasibility, Bakken et al. (2012) evaluated the technical requirements and constraints for offshore groundwater abstraction, transport, and treatment, concluding that OFG could serve as a viable drinking water source without significant technical, economic, or environmental impediments.

Potential adverse impacts associated with the utilisation of OFG include reduced availability of onshore groundwater and subsidence (Yu and Michael 2019), loss or contamination of submarine groundwater discharge affecting ecosystems (Varma and Michael 2012) and seawater intrusion in coastal aquifers (Post and Houben 2017). Additional concerns involve brine disposal, marine habitat disturbance and drilling-related pollution (Ghaffour et al. 2013).

In parallel to technical and environmental considerations, the broader sustainability and societal dimensions of OFG development must be assessed. Early stakeholder consultations by the OFF-SOURCE network highlighted widespread concern that OFG development could replicate past patterns of over-abstraction seen in many coastal aquifers, especially where institutional capacity is limited. A consistent message was that OFG should not be pursued as an isolated solution but recognised as a component of an integrated water resource management strategy, particularly where population growth as well as climatic variability and hydrological extremes are intensifying groundwater demand and reducing reliability of traditional sources.

An OFF-SOURCE report identified four potential utilisation scenarios for OFG: (a) direct use with minimal treatment for non-potable applications; (b) reverse osmosis treatment to meet full potable water standards; (c) strategic reserve accessed only during supply emergencies; and (d) offshore use without land transport to support marine industrial operations (OFF-SOURCE 2024). Stakeholder consultations emphasized that realizing these scenarios requires building synergies with existing offshore industries—including oil and gas, wind energy, and carbon storage—through shared infrastructure, coordinated economic models, and aligned regulatory frameworks.

Legal challenges and ownership

As scientific evidence of vast quantities of OFG grows and demand of freshwater for coastal areas increases, both sovereign states and private entities will begin to contemplate accessing OFG to supplement or even replace dwindling onshore reserves of freshwater. While solutions to the technical and financial hurdles are being explored, questions of ownership and governance on OFG resources will naturally arise.

As of 2025, 171 states have ratified (or acceded to) the United Nations (UN) Convention on the Law of the Sea (UNCLOS), meaning that the Convention offers a nearly universal framework for allocating sovereign rights over marine resources. Pursuant to UNCLOS, coastal States have full sovereignty over the Territorial Sea, extending up to twelve nautical miles from the low-tide line, including the water column, seabed, and subseafloor (Figure S1 of the ESM). Beyond the Territorial Sea and for two hundred miles from the low tide line, in an area called the Exclusive Economic Zone (EEZ), nations have exclusive sovereign rights to all natural resources. The Territorial Sea and the EEZ together are often called the continental shelf, although in some cases the continental shelf does not extend for the full length of the EEZ, and in other cases the continental shelf extends further than the EEZ, which gives the UNCLOS Member State additional rights. Since deposits of OFG will be located within the continental shelf, the right of a coastal nation to explore and exploit the freshened water lying exclusively within its marine territory will not be disputed, and nations will establish their own laws and regulations regarding exploration, extraction, transport, and use of the resource.

However, natural resources do not respect political boundaries, and some deposits of OFG will straddle the continental shelf of one or more nations. Since UNCLOS provides no guidance on transboundary seabed resources, nations will seek guidance when crafting a framework for ownership and governance. OFG is similar to O&G on the marine subsoil, in that the deposits are contained but fluid,

so reference to offshore O&G arrangements would be logical. In addition, multiple treaties exist to allocate rights and responsibilities for transboundary offshore resources, especially valuable reserves of O&G, so those agreements could serve as precedents. The most common method of allocating costs and benefits of finding and extracting offshore O&G is known as a joint development agreement (JDA), wherein nations agree in advance to share the costs and revenues of offshore reserves in accordance with the proportional share of the reserve that each of them is assumed to hold. For efficiency and to reduce costs, the parties also agree to appoint one entity to represent them both in the exploration and development activities.

In addition to questions regarding ownership of OFG, nations will also have to consider the environmental impacts of OFG development, such as the effects of operational noise, pollution, and sediment disruption on marine species as well as the potential for seafloor subsidence. The environmental concerns can become more complicated when the impacts are transboundary. Certain global treaties, such as the Convention on Biological Diversity and the Espoo Convention, may apply, and OFG development will also have to conform to domestic environmental laws and regulations that address marine activities and freshwater quality and distribution.

Within the European context, a key water legislation is the Water Framework Directive (WFD), which establishes a framework for the protection and sustainable management of surface waters and groundwater. When implemented in conjunction with international water treaties and domestic laws, the WFD can serve as a legally binding instrument to prevent water deterioration and preserve groundwater dependent ecosystems. However, the current definition of groundwater in (Article 2, para 2) of the WFD, “water below the surface of the ground in the saturation zone and in direct contact with the ground or subsoil” raises interpretative questions. The wording ‘direct contact with the ground or subsoil’ may be understood as excluding ‘fossil’ groundwater that is disconnected from the land surface. Aquifer is further defined as “subsurface layer or layers of rock or other geological strata of sufficient porosity and permeability to allow either a significant flow of groundwater or the abstraction of significant quantities of groundwater” (Article 2, para 11). It then follows that deep aquifers that are not used for groundwater abstraction need not be identified as groundwater bodies. Moreover, it remains unclear whether the terms “subsoil” or “the ground” encompass the seabed and its subsoil, thereby creating uncertainty for the regulation of offshore fresh groundwater.

Furthermore, the Marine Strategy Framework Directive should be revised to ensure that changes in SGD quality and discharge volumes resulting from OFG utilisation are incorporated as pressures that may affect the achievement of good

environmental status. Additionally, marine spatial planning (MSP), which is part of the Integrated Maritime Policy of the EU, also lacks mention of OFG. Bringing this topic early into MSP is essential to both give a planning aspect to OFG utilisation, and safeguard other activities from potential OFG utilisation risks. Thus, coastal nations contemplating development of OFG would be wise to plan how to address these various governance and environmental issues.

Summary

The science of OFG has significantly advanced in the last decades, thanks to improvement in research tools, data acquisition and processing, and modelling techniques. However, a series of questions are still open, mainly related to OFG recharge mechanisms, total volumes, connectivity with onshore coastal aquifers, its response to sea-level and climate variability and its influence on marine ecosystems.

Sustained action to consolidate data from various sectors and facilitate collaborative research, knowledge exchange, and policy formulation, are needed to ensure a harmonised approach to OFG research, and to its management and utilisation worldwide.

Continued development of integrated geological and geophysical methods and 3D numerical modelling is needed to provide more robust spatial and volumetric analysis of OFG systems. This can be facilitated by testing theory-guided ML frameworks. ML-enabled database generation (such as from raster digitisation of O&G legacy well data) has also the potential to offer a practical means of mitigating the chronic data scarcity that hampers OFG research.

In terms of utilisation, while some factors, such as transport logistics and local infrastructure, are geographically specific, others, like basic resource availability and treatment technologies, have broader applicability across various regions. OFG should be recognised as one component of an integrated onshore-offshore coastal water-resources strategy—particularly where population growth as well as extreme hydroclimatic variability is intensifying groundwater demand and reducing reliability of traditional sources. This will depend on both global and local factors.

Finally, coastal nations contemplating development of OFG should plan how to address the various governance and environmental issues, drawing from existing offshore resources treaties and regulations. At the European level, the WFD should be modified to cover deep groundwater, which is disconnected from surface systems, both on land and offshore, while the Marine Strategy Framework Directive should be revised to incorporate changes in SGD quality and discharge volumes resulting from OFG utilisation.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10040-026-03054-x>.

Acknowledgements This article/publication is based upon work from CA21112 – “Offshore freshened groundwater: An unconventional water resource in coastal regions? – OFF-SOURCE”, supported by COST (European Cooperation in Science and Technology). CB, AC, MG, EQJ, IS would like to thank the European Commission and the Water4All Partnership for funding in the framework of the collaborative international consortium RESCUE financed under the 2022 Joint call of the European Partnership 101060874 — Water4All. D Gamboa thanks financial support by national funds from FCT – Fundação para a Ciência e Tecnologia I.P, to project CESAM-Centro de Estudos do Ambiente e do Mar, references UID/50017/2025 (<https://doi.org/10.54499/UID/50017/2025>), LA/P/0094/2020 (<https://doi.org/10.54499/LA/P/0094/2020>). AM was supported by the David and Lucile Packard Foundation. The authors are grateful to Vincent Post for his thoughtful review of the manuscript and helpful suggestions.

Author contributions All authors contributed to the study conception and design. CB, AT and HW provided the first structure and content table of the manuscript, and AC, JCC, MG, DG, EQJ, RMN, AM, MN and IS wrote relevant sections of the manuscript on previous versions of the manuscript. All authors read and approved the final manuscript.”

Funding Open access funding provided by Università degli Studi di Trieste within the CRUI-CARE Agreement. CA21112 – “Offshore freshened groundwater: An unconventional water resource in coastal regions? – OFF-SOURCE”, supported by COST (European Cooperation in Science and Technology). RESCUE: European Commission 2022 Joint call of the European Partnership 101060874 — Water4All.

Declarations

Competing interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Adombi AVDP, Boucher M-A, Chesnaux R (2021) Review: theory-guided machine learning applied to hydrogeology—state of the art, opportunities and future challenges. *Hydrogeol J* 29(8):2671–2683. <https://doi.org/10.1007/s10040-021-02403-2>
- Arévalo-Martínez DL, Haroon A, Bange HW, Erkul E, Jegen M, Moosdorf N et al (2023) Ideas and perspectives: land–ocean connectivity through groundwater. *Biogeosciences* 20(3):647–662. <https://doi.org/10.5194/bg-20-647-2023>
- Bakken TH, Ruden F, Mangset LE (2012) Submarine groundwater: a new concept for the supply of drinking water. *Water Resour Manag* 26(4):1015–1026. <https://doi.org/10.1007/s11269-011-9806-1>
- Bertoni C, Lofi J, Micallef A, Moe H (2020) Seismic reflection methods in offshore groundwater research. *Geosciences* 10(8):299. <https://doi.org/10.3390/geosciences10080299>
- Campo B, Pellegrini C, Sammartino I, Trincardi F, Amorosi A (2024) New perspectives on offshore groundwater exploration through integrated sequence-stratigraphy and source-to-sink analysis: insights from the late Quaternary succession of the western Central Adriatic system, Italy. *Earth-Sci Rev* 256:104880. <https://doi.org/10.1016/j.earscirev.2024.104880>
- Chiacchieri D, Lipparini L, Jordan EQ, Bencini R, Micallef A (2025) Coastal fresh groundwater extending deep offshore Southern Sicily (Italy): assessment of the ragusa aquifer via petrophysical and 3D hydrogeological modelling. *Hydro J*(10)
- Corradi C, Thomas AT, Camerlenghi A, Zini L, Giustiniani M, Busetti M, Foglia L, Bertoni C, Micallef A (2025) Characterization of an onshore-offshore aquifer system in the Venetian Friulian Plain and north-eastern Adriatic. *Hydro J* (10)
- Cracknell MJ, Reading AM (2014) Geological mapping using remote sensing data: a comparison of five machine learning algorithms, their response to variations in the spatial distribution of training data and the use of explicit spatial information. *Comput Geosci* 63:22–33. <https://doi.org/10.1016/j.cageo.2013.10.008>
- De Biase M, Chidichimo F, Micallef A, Cohen D, Gable C, Zwinger T (2023) Past and future evolution of the onshore-offshore groundwater system of a carbonate archipelago: the case of the Maltese Islands, central Mediterranean Sea. *Front Water* 4:1068971. <https://doi.org/10.3389/frwa.2022.1068971>
- Faghih Z, Haroon A, Jegen M, Gehrmann R, Schwalenberg K, Micallef A et al (2024) Characterizing offshore freshened groundwater salinity patterns using trans-dimensional Bayesian inversion of controlled source electromagnetic data: a case study from the Canterbury Bight, New Zealand. *Water Resour Res* 60(3):e2023WR035714. <https://doi.org/10.1029/2023WR035714>
- Ghaffour N, Missimer TM, Amy GL (2013) Technical review and evaluation of the economics of water desalination: current and future challenges for better water supply sustainability. *Desalination* 309:197–207. <https://doi.org/10.1016/j.desal.2012.10.015>
- Giustiniani M, Gamboa D, Bertoni C, Usmand N, Thomas AT, Micallef A, Retike I, Bachtouli S, Barison E, Busetti M, Corradin C, Garcia X, Leon R, Lobo F, Hermans T, Hoffmann J, Lofi J, Paepen M, Pambuku A, Rossi V, Shaban A, Tziritis E, Vignoli G, Zairi M, Wazaz HA (2025) Geological and Geophysical database for the assessment of Offshore Freshened Groundwater resources. *Hydro J* (10)
- Gupta S, Micallef A (2025) Numerical modelling of erosional landforms driven by offshore groundwater flow on siliciclastic continental margins: a conceptual approach. *Front Earth Sci* 13:1453255. <https://doi.org/10.3389/feart.2025.1453255>
- Haffert L, Jegen M, Siebert C, Rödiger T, Berndt C (2024) AI-Quifer - using artificial intelligence to determine offshore groundwater occurrences that are key to coastal water management. In: OCEANS. Halifax, pp 1–5
- Hathaway JC, Poag CW, Valent PC, Miller RE, Schultz DM, Manhe FT, et al (1979) US Geological Survey Core Drilling on the Atlantic Shelf. In *Science* 206 (4418), pp 515–527
- Hoffmann JLL, Mountjoy JJ, Spain E, Gall M, Tait LW, Ladroit Y, Micallef A (2023) Fresh submarine groundwater discharge offshore Wellington (New Zealand): hydroacoustic characteristics and its influence on seafloor geomorphology. *Front Mar Sci* 10:1204182. <https://doi.org/10.3389/fmars.2023.1204182>
- Horozal S, Micallef A, De Biase M, Chidichimo F (2025) Offshore freshened groundwater in an active rift: Gulf of Corinth, Greece. *Hydro J* (10)
- Ishizu K, Ogawa Y (2021) Offshore-onshore resistivity imaging of freshwater using a controlled-source electromagnetic method: a feasibility study. *Geophysics* 86(6):E391–E405, -. <https://doi.org/10.1190/geo2020-0906.1>
- IODP³ (2025) New England Shelf Hydrogeology (Joint IODP3-NSF Expedition), see: <https://iodp3.org/expedition/501>. Accessed 23 February 2026

- JPI Oceans (2025) Offshore freshened groundwater. See: <https://www.jpi-oceans.eu/en/offshore-freshened-groundwater>. Accessed 23 February 2026
- Katole AL, Mangsuli P, Gune OA (2025) Rejuvenating legacy data by digitizing raster logs. In *Artificial Intelligence for Subsurface Characterization and Monitoring*, p. 3–22, Elsevier, Netherlands, 269pp
- Knight AC, Werner AD, Morgan LK (2018) The onshore influence of offshore fresh groundwater. *J Hydrol* 561:724–736. <https://doi.org/10.1016/j.jhydrol.2018.03.028>
- Kohout FA (1964) The flow of fresh water and salt water in the Biscayne Bay aquifer of the Miami area, Florida. *Sea water in coastal aquifers*. In *US. Geol. Surv., Water Supply Pap.*, pp. 12–32. Available online at <https://cir.nii.ac.jp/crid/1571698599164155648>
- Li F, Li M, Comte J-C, Wu P, Chen H (2025) Hydrological characteristics of Nearshore Freshened Groundwater (NFG) and its environmental effects of exploitation in the east coast of Laizhou Bay, China. *Hydro J* (10)
- Lin X, Li J, Zhang T, Ma Q, Abdullah A (2025) Aquitard brine evolution from paleoseawater during the Late Pleistocene in the coastal plain of the southern Laizhou Bay (China): hydrochemical and isotopic evidences. *Hydro J* (10)
- Lipparini L, Chiacchieri D, Bencini R, Micallef A (2023) Extensive freshened groundwater resources emplaced during the Messinian sea-level drawdown in southern Sicily, Italy. *Commun Earth Environ* 4(1):430. <https://doi.org/10.1038/s43247-023-01077-w>
- Martin-Nagle R (2020) Offshore freshwater aquifers: toward equitable distribution. *Wires Water* 7(5):e1463. <https://doi.org/10.1002/wat2.1463>
- Micallef A, Person M, Berndt C, Bertoni C, Cohen D, Dugan B et al (2021) Offshore freshened groundwater in continental margins. *Rev Geophys*. <https://doi.org/10.1029/2020RG000706>
- Micallef A, Person M, Gupta S, Saadatkah N, Camille A, Gratacós Ò (2023) Can offshore meteoric groundwater generate mechanical instabilities in passive continental margins? *J Geophys Res Earth Surf* 128:e2022JF006954
- Moore WS (2010) The effect of submarine groundwater discharge on the ocean. In *Annual review of marine science* 2 (Volume 2, 2010), pp 59–88. <https://doi.org/10.1146/annurev-marine-120308-081019>
- NDR repository (n.d.) <https://ndr.nstauthority.co.uk/>. Accessed 13 Mar 2026
- OFF-SOURCE (2024). Unlocking Offshore Freshened Groundwater. OFF-SOURCE COST Action (CA21112). 10.5281/zenodo.17804632
- Pastorella AE, Haroon A, Everett ME, Rohde L, Bartels T, Wollatz-Vogt M, Faghih Z, Franz GK, Micallef A (2023) SWAN: a surface-towed modular controlled-source electromagnetic system for mapping submarine groundwater discharge and offshore groundwater resources. *Lead Edge* 42:590–597
- Post VEA, Groen J, Kooi H, Person M, Ge S, Edmunds WM (2013) Offshore fresh groundwater reserves as a global phenomenon. *Nature* 504(7478):71–78. <https://doi.org/10.1038/nature12858>
- Post VEA, Houben GJ (2017) Density-driven vertical transport of saltwater through the freshwater lens on the island of Baltrum (Germany) following the 1962 storm flood. *J Hydrol* 551:689–702. <https://doi.org/10.1016/j.jhydrol.2017.02.007>
- Pourghasemi HR, Sadhasivam N, Yousefi S, Tavangar S, Ghaffari Nazarlou H, Santosh M (2020) Using machine learning algorithms to map the groundwater recharge potential zones. *J Environ Manage* 265:110525. <https://doi.org/10.1016/j.jenvman.2020.110525>
- RESCUE (2026) Resources in coastal groundwater under hydroclimatic extremes, see <https://www.rescue-water4all.eu/>. Accessed 23 February 2026.
- Quiroga E, Bertoni C, Ruden F (2023) Deep low-salinity groundwater in sedimentary basins: petrophysical methods from a case study in Somalia. *Hydrogeol J* 31(3):685–705. <https://doi.org/10.1007/s10040-022-02589-z>
- Rahman et al. (2025) Deep fresh groundwater in offshore islands and its implications for climate-resilient groundwater management in coastal aquifers. *Hydro J* (10)
- Roberts MJ, Vidale PL, Senior C, Hewitt HT, Bates C, Berthou S et al (2018) The benefits of global high resolution for climate simulation: process understanding and the enabling of stakeholder decisions at the regional scale. *Bull Am Meteorol Soc* 99(11):2341–2359. <https://doi.org/10.1175/bams-d-15-00320.1>
- Ruden F (2007) The discovery of a regional Neogene aquifer in coastal Tanzania. *Coastal Aquifers: Challenges and Solutions* 1:363–372
- Saadatkah N, Kassim A, Siat QA, Micallef A (2023) Salt leaching by freshwater and its impact on seafloor stability: an experimental investigation. *Mar Geol* 455:106959. <https://doi.org/10.1016/j.margeo.2022.106959>
- Santos IR, Burnett WC, Chanton J, Mwashote B, Suryaputra IGNA, Dittmar T (2008) Nutrient biogeochemistry in a Gulf of Mexico subterranean estuary and groundwater-derived fluxes to the coastal ocean. *Limnol Oceanogr* 53(2):705–718. <https://doi.org/10.4319/lo.2008.53.2.0705>
- Sheng C, Jiao JJ, Luo X, Zuo J, Jia L, Cao J (2023) Offshore freshened groundwater in the Pearl River estuary and shelf as a significant water resource. *Nat Commun* 14(1):3781. <https://doi.org/10.1038/s41467-023-39507-0>
- Sola F, Fernandez-Cortes A, Vallejos A (2025) Paleo-coastline reconstruction based on the hydrogeochemistry of coastal aquifers. *Hydro J* (10)
- Taniguchi M, Burnett WC, Cable JE, Turner JV (2002) Investigation of submarine groundwater discharge. *Hydrol Process* 16(11):2115–2129. <https://doi.org/10.1002/hyp.1145>
- Thomas AT, Harten JV, Jusri T, Reiche S, Wellmann F (2022) An integrated modeling scheme for characterizing 3D hydrogeological heterogeneity of the New Jersey shelf. *Mar Geophys Res* 43(2):11. <https://doi.org/10.1007/s11001-022-09475-z>
- Thomas AT, Zamrsky D, Oude E, Gualbert HP, Bierkens MFP, Micallef A (2025) Offshore Freshened Groundwater prospecting using Machine Learning.
- UN-Water (2020) UN-Water Analytical Brief on Unconventional Water Resources. Geneva, Switzerland. Available online at <https://www.unwater.org/publications/un-water-analytical-brief-on-unconventional-water-resources/>
- Varma S, Michael K (2012) Impact of multi-purpose aquifer utilisation on a variable-density groundwater flow system in the Gippsland Basin, Australia. *Hydrogeol J* 20(1):119–134
- VIDEPI repository (n.d.) <https://www.videpi.com/videpi/videpi.asp>. Accessed 13 Mar 2026
- Waliser D, Seo K-W, Schubert S, Njoku E (2007) Global water cycle agreement in the climate models assessed in the IPCC AR4. *Geophys Res Lett* 34(16):2007GL030675. <https://doi.org/10.1029/2007GL030675>
- Water Framework Directive (2000) Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy. *WFD L 327*, pp. 1–73. Available online at [https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32000L0060&from=fr#:~:text=\(f\)%20controls%2C%20including%20a,provided%20for%20under%20Community%20legislation](https://eur-lex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32000L0060&from=fr#:~:text=(f)%20controls%2C%20including%20a,provided%20for%20under%20Community%20legislation)
- Yu S, Kurylyk BL, Michael HA, Maselli V, Nedimović MR, Schulten I, Córdoba-Ramírez F (2025) Numerical investigation of offshore freshened groundwater dynamics in a changing marine environment. *J Hydrol* 664:134553. <https://doi.org/10.1016/j.jhydrol.2025.134553>
- Yu X, Michael HA (2019) Offshore pumping impacts onshore groundwater resources and land subsidence. *Geophys Res Lett* 46(5):2553–2562. <https://doi.org/10.1029/2019GL081910>
- Zamrsky D, Essink GHPO, Sutanudjaja EH, van Beek LPH, Bierkens MFP (2022) Offshore fresh groundwater in coastal unconsolidated sediment systems as a potential fresh water source in the 21st century. *Environ Res Lett* 17(1):14021. <https://doi.org/10.1088/1748-9326/ac4073>